CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 104, 2011

A

Abe, R., 646
Abe, Y., 1771
Abeyo, B. G., 2073
Abidha, S., 1816
Abney, M. R., 143
Abolmaaty, A., 41
Adán, A., 1541
Agnello, A. M., 1613
Aguado, P., 1541
Akbulut, M., 1212
Albajes, R., 853
Alexander, L. C., 1316
Alfaro, C., 1647
Alim, M. A., 1833
Allison, J. D., 895, 1258
Alm, S. R., 41
Almeida, R.P.P., 1480
Aluja, M., 1204
Alzogaray, R. A., 595
An, K., 1840
Anciso, J., 1138
Anderson, K. M., 1055, 1393
Andreller, I. S., 1018
Anfora, G., 1125
Anulewicz, A. C., 517, 1599
Arbogast, R. T., 26, 1430
Arévalo-Galarza, L., 75
Arieli, B., 1633
Armstrong, J. W., 1509
Arouri, R., 1349
Arrington, M., 137
Arthur, F. H., 702, 1415, 1765
Artz, D. R., 1153
Asaro, Christopher, 1245, 1272
Athanassiou, C. G., 1752, 1765
Attia, S., 1220
Avelino, N., 1018
Ayer, T., 379
Ayvaz, A., 1212

B

Baenziger, P. S., 2073
Bagley, M., 1038
Bai, Y., 1892
Baker, P. B., 1339
Ball, O. J.-P., 665
Ballman, E. S., 279
Barak, A. V., 69
Barbour, a.J.D., 933
Barden, S. A., 2009
Barr, N. B., 920, 1706
Barraco, G., 180
Barrington, A. M., 1301
Bartelt, R. J., 1195
Bastos, C. S., 1909
Bauer, L. S., 81
Baute, T. S., 343
Baxendale, F. P., 205
Beauzay, P. B., 1236
Beeby, R. L., 308
Belay, D. K., 1329
Bellini, R., 266
Bengochea, P., 1541
Berenbaum, M. R., 459
Beroiz, B., 1349
Bharadwaj, A., 2095
Birke, A., 1204
Birmingham, A. L., 1018
Björkman, C., 782
Blankenship, E. E., 205
Blomefield, T., 814
Boettner, G., 494
Bommarco, R., 782
Boncristiani, H., 732
Bonjour, E. L., 308, 1415
Borden, J. H., 1018
Bosch, J., 750
Bosco, D., 360, 1480
Bosque-Pérez, N. A., 164
Botch, P. S., 425
Boudreau, E., 1584
Bowen, K. I., 1167
Bradshaw, J. D., 1561
Braman, S. K., 288, 868, 1918
Brockerhoff, E. G., 1514
Bu, C. Y., 375
Buchman, J. L., 1486, 1783
Budia, F., 1229, 1541
Bukowsky, R., 2054
Bullangpoti, V., 1534
Burkness, E. C., 1476, 1900
Burks, C. S., 211
Buschman, L. L., 343
Butler, C. D., 586
Butler, R. C., 1999
Butler, S. M., 1258
Bylund, H., 782
Byrne, A., 1357
Byrne, F. J., 586, 868

C

Cambron, S., 1055
Campbell, J. F., 443, 2078
Campbell, L., 2068
Campos, S. O., 1909
Cañas, L., 548, 574
Cappaert, D., 1599
Cárcamo, H. A., 1550
Cardona, P. S., 1406
Carloni, E., 1793
Carpane, P., 1793
Carpenter, J. E., 814
Carraher, C., 920
Carrière, Y., 1339
Carrieri, M., 266
Carvalho, S. M., 636
Castañera, P., 1349
Castle, S. J., 773
Catal, M., 839
Cervera, A., 1349
Chandrapatya, A., 1445
Chandrasena, D., 1357
Chang, Z., 1502
Charlet, L. D., 1236
Chaudhury, M. F., 1103
Chediak, M., 1909
Chen, H., 1730
Chen, J., 1877
Chen, J.-L., 1080
Chen, Li, 1436, 1877
Chen, Liezhong, 1877
Chen, Lizhen, 1189
Chen, M., 709
Chen, M.-S., 1384, 2068
Chen, W.-T., 225
Chen, X. M., 149
Chen, Yanping, 732
Chen, Y.-C., 614
Chen, Yi, 1592
Chen, Yigen, 81
Chen, You, 149
Chen, Y.-Y., 225
Cheng, C.-C., 823
Cheng, D.-F., 1080
Cheng, J., 375, 1892
Cherry, R. H., 155, 474
Chevrier, C., 188
Chi, W.-L., 225
Chirumamilla, A., 1236
Chu, D., 978
Chun, Y. S., 693
Ciaramitaro, T., 173
Cloyd, R. A., 1633
Cohnstaedt, L. W., 1430
Conway, H. E., 452
Cooley, D. R., 1613
Cooper, R., 274
Cooper, W. R., 685
Cornelius, M. L., 1009, 1024
Cottrell, T. E., 14, 47
Coudron, T. A., 414
Couso-Ferrer, F., 1349
Creighton, J., 1272
Cristofaro, A. D., 1125
Crowe, C. M., 1245, 1934
Cuda, J. P., 1986

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 104, 2011

A

Abe, R., 646
Abe, Y., 1771
Abeyo, B. G., 2073
Abidha, S., 1816
Abney, M. R., 143
Abolmaaty, A., 41
Adán, A., 1541
Agnello, A. M., 1613
Aguado, P., 1541
Akbulut, M., 1212
Albajes, R., 853
Alexander, L. C., 1316
Alfaro, C., 1647
Alim, M. A., 1833
Allison, J. D., 895, 1258
Alm, S. R., 41
Almeida, R.P.P., 1480
Aluja, M., 1204
Alzogaray, R. A., 595
An, K., 1840
Anciso, J., 1138
Anderson, K. M., 1055, 1393
Andreller, I. S., 1018
Anfora, G., 1125
Anulewicz, A. C., 517, 1599
Arbogast, R. T., 26, 1430
Arévalo-Galarza, L., 75
Arieli, B., 1633
Armstrong, J. W., 1509
Arouri, R., 1349
Arrington, M., 137
Arthur, F. H., 702, 1415, 1765
Artz, D. R., 1153
Asaro, Christopher, 1245, 1272
Athanassiou, C. G., 1752, 1765
Attia, S., 1220
Avelino, N., 1018
Ayer, T., 379
Ayvaz, A., 1212

B

Baenziger, P. S., 2073
Bagley, M., 1038
Bai, Y., 1892
Baker, P. B., 1339
Ball, O. J.-P., 665
Ballman, E. S., 279
Barak, A. V., 69
Barbour, a.J.D., 933
Barden, S. A., 2009
Barr, N. B., 920, 1706
Barraco, G., 180
Barrington, A. M., 1301
Bartelt, R. J., 1195
Bastos, C. S., 1909
Bauer, L. S., 81
Baute, T. S., 343
Baxendale, F. P., 205
Beauzay, P. B., 1236
Beeby, R. L., 308
Belay, D. K., 1329
Bellini, R., 266
Bengochea, P., 1541
Berenbaum, M. R., 459
Beroiz, B., 1349
Bharadwaj, A., 2095
Birke, A., 1204
Birmingham, A. L., 1018
Björkman, C., 782
Blankenship, E. E., 205
Blomefield, T., 814
Boettner, G., 494
Bommarco, R., 782
Boncristiani, H., 732
Bonjour, E. L., 308, 1415
Borden, J. H., 1018
Bosch, J., 750
Bosco, D., 360, 1480
Bosque-Pérez, N. A., 164
Botch, P. S., 425
Boudreau, E., 1584
Bowen, K. I., 1167
Bradshaw, J. D., 1561
Braman, S. K., 288, 868, 1918
Brockerhoff, E. G., 1514
Bu, C. Y., 375
Buchman, J. L., 1486, 1783
Budia, F., 1229, 1541
Bukowsky, R., 2054
Bullangpoti, V., 1534
Burkness, E. C., 1476, 1900
Burks, C. S., 211
Buschman, L. L., 343
Butler, C. D., 586
Butler, R. C., 1999
Butler, S. M., 1258
Bylund, H., 782
Byrne, A., 1357
Byrne, F. J., 586, 868

C

Cambron, S., 1055
Campbell, J. F., 443, 2078
Campbell, L., 2068
Campos, S. O., 1909
Cañas, L., 548, 574
Cappaert, D., 1599
Cárcamo, H. A., 1550
Cardona, P. S., 1406
Carloni, E., 1793
Carpane, P., 1793
Carpenter, J. E., 814
Carraher, C., 920
Carrière, Y., 1339
Carrieri, M., 266
Carvalho, S. M., 636
Castañera, P., 1349
Castle, S. J., 773
Catal, M., 839
Cervera, A., 1349
Chandrapatya, A., 1445
Chandrasena, D., 1357
Chang, Z., 1502
Charlet, L. D., 1236
Chaudhury, M. F., 1103
Chediak, M., 1909
Chen, H., 1730
Chen, J., 1877
Chen, J.-L., 1080
Chen, Li, 1436, 1877
Chen, Liezhong, 1877
Chen, Lizhen, 1189
Chen, M., 709
Chen, M.-S., 1384, 2068
Chen, W.-T., 225
Chen, X. M., 149
Chen, Yanping, 732
Chen, Y.-C., 614
Chen, Yi, 1592
Chen, Yigen, 81
Chen, You, 149
Chen, Y.-Y., 225
Cheng, C.-C., 823
Cheng, D.-F., 1080
Cheng, J., 375, 1892
Cherry, R. H., 155, 474
Chevrier, C., 188
Chi, W.-L., 225
Chirumamilla, A., 1236
Chu, D., 978
Chun, Y. S., 693
Ciaramitaro, T., 173
Cloyd, R. A., 1633
Cohnstaedt, L. W., 1430
Conway, H. E., 452
Cooley, D. R., 1613
Cooper, R., 274
Cooper, W. R., 685
Cornelius, M. L., 1009, 1024
Cottrell, T. E., 14, 47
Coudron, T. A., 414
Couso-Ferrer, F., 1349
Creighton, J., 1272
Cristofaro, A. D., 1125
Crowe, C. M., 1245, 1934
Cuda, J. P., 1986

Cui, J., 673
Culler, L. E., 1316
Cunningham, N. M., 220
Currie, R. W., 1774

D
Daane, K. M., 367, 1480
Dai, F.-Y., 101
Daneel, J. H., 1174, 1180
Danka, R. G., 745
Delucia, E. H., 459
Demichelis, S., 360
Demir, H., 839
Deng, T., 862
Diaz-Montano, J., 1
Diaz-Sobac, R., 1991
Dickinson, A., 1633
Dierig, D. A., 1575
Difonzo, C., 343, 1357
Dindo, M. L., 266
Dodds, K. J., 879
Dogan, H., 325
Dolan, M. C., 659, 2062
Dosdall, L. M., 1550
Dou, W., 1436, 1720
Dowell, R. V., 1514
Dreves, A. J., 970
Duan, D. D., 375
Duan, J. J., 81
Duehl, A. J., 1430
Duerr, D. A., 1245
Duraimurugan, P., 1406
Dyson, C., 220

E
Eaton, M., 1423
Ecroyd, C. E., 1514
Edu, J., 1969
Eigenbrode, S. D., 164
Eizaguirre, M., 853
Ekesi, S., 1180
Elder, P., 69
Elkinton, J. S., 494
Ellersieck, M., 1045, 1584
Ellsworth, P. C., 1575
El-Sayed, A. M., 1301
El-Zemaity, M. E.-S., 41
England, K. M., 548
Enkegaard, A., 799
Epstein, M., 920, 1706
Erler, F., 839
Estal, P. D., 1541

F
Fadhl, S., 188
Fail, J., 1
Fang, C.-C., 2039
Fang, S.-M., 101
Farris, R. E., 1706
Fathipour, Y., 799
Faubert, H., 1613
Feng, H.-T., 2039
Fernandes, F. L., 1909
Fields, P. G., 1445
Figueroa, C. C., 258
Figueroa, J.-I., 1229
Flanders, K. L., 1167
Flinn, P. W., 317, 325
Flores, S., 1885
Folgarait, P. J., 32
Follett, P. A., 75, 566, 571, 1509
Foo, F.-K., 1675
Forbes, A. A., 331
Forrester, O. T., 452
Foster, J. E., 1329
Frake, A. M., 26, 1146
Francese, J. A., 901
Francis, a. F., 1080
Frank, D. L., 1584, 2054
Frank, S. D., 1960
Fraser, I., 69, 379, 901
French, B. W., 1038, 2054
Frost, K. E., 1800
Fuchs, M., 1
Furlong, M. J., 629

G
Gadino, A. N., 970
Galetto, L., 360
Gandhi, K.J.K., 574
Ganehiarachchi, G.a.s.m., 1236
Ganesan, K., 1369
Gao, C., 1364
Gao, L. W., 965
Gao, Y., 1771
Gardescu, S., 1942
Garud, A., 1369
Gaussoin, R. E., 205
Gautam, B. K., 2025
Ge, D., 978
Geden, C. J., 1108
Gerry, A. C., 1093
Gilbert, R. A., 155, 474
Gill, T. A., 299
Gilligan, T. M., 920, 1706
Godfrey, A.B.P.L.D., 1294
Gomez, L. E., 1856, 1885
Goolsby, J., 1138
Gould, J. R., 379
Goulson, D., 107
Gradziel, T., 965
Grafton-Cardwell, E. E., 1286
Graham, L. C., 2009
Grant, G. G., 173
Grant, J. A., 1496
Gratton, C., 115, 1568
Gray, M. E., 1561
Gregory, A., 1138
Grenkow, L. F., 125
Grgurinovic, C. A., 717
Gries, G., 1018
Grissa, K. L., 1220
Gross, K., 331
Grout, T. G., 1174, 1180
Groves, R. L., 1800
Gruber, M. Y., 125
Guedes, R.n.c., 1909
Guerfali, M. M., 188
Guillade, A. C., 32
Guo, K., 1851
Guo, Y., 827
Guzman, L.I.D., 26, 1146
Gwinn, K. D., 665
Gwirtz, J. A., 317, 325

H
Haack, R. A., 717
Hadi, B.A.R., 1167
Hajek, A. E., 1942
Halbert, S. E., 1167
Hallett, R. H., 709
Hallman, G. J., 21, 63, 1851
Hama, M., 646
Hamasaki, R. T., 566
Hamby, K., 965
Hamden, H., 188
Hammond, A., 1138
Hance, T., 1220
Hanks, L. M., 933, 1592
Hanula, J. L., 288, 501, 868, 1918
Haraguchi, D., 420
Hardin, J., 308
Harrand, L., 1087
Harris, J. W., 745
Harris, M. O., 1055, 1393
Hashiyama, A., 1280
Hassan, M. W., 1436
Hassanpour, M., 1379
Hattingh, V., 1174, 1180
He, Yueping, 1364, 1877
He, Yuxian, 833
Headrick, D. H., 1286
Heilman, B. E., 1783
Hein, G. L., 1900
Held, D. W., 2009
Hellmich, R. L., 343
Henderson, G., 1002, 2025
Heng-Moss, T. M., 205, 2073
Herbert, Jr., D. A., 137
Herms, D. A., 574
Hernández-Crespo, P., 1349
Heuskin, S., 1220
Hibbard, B. E., 1045, 1584, 2054
Higbee, B. S., 211
Higdon, M. L., 1045
Higginbotham, G. E., 1093
Hilje, L., 164
Hirase, K., 646
Hlasny, V., 764
Ho, H.-Y., 823
Hoddle, M. S., 279
Hoepting, C. A., 709
Hoffmann, A. A., 465
Hogg, D. B., 115
Hogsette, J. A., 1108
Holliday, N. J., 889

Holman, D., 1138
Holowachuk, J. M., 125
Honěk, A., 1870
Hong, S. C., 1568
Hooks, C.R.R., 947
Hoover, K., 717
Horn, S., 501
Horton, D. L., 47
Hsu, J.-C., 2039
Huang, F., 1892
Huang, Jia, 1840, 2046
Huang, Z. Y., 149
Hulbert, D., 1656
Hummer, K., 566
Hunt, T. E., 1900
Hurley, T. M., 343
Hussien, M. I., 41
Hutchison, W. D., 343, 1476, 1900
Hwang, S.-Y., 1525

I

Iii, K.C.S., 2095
Ioriatti, C., 1125
Isaacs, R., 1656
Ishtiaq, M., 1343
Islam, A., 21

J

Jalaludin, N. A., 607
James, R. R., 353
Jankielsohn, A., 1736
Jaramillo, J., 87
Jarosík, V., 1870
Jennings, S. N., 107
Jessup, A. J., 21
Ji, J., 693
Jiang, H.-B., 1436
Jiao, X., 978
Johnson, C. W., 1258
Johnson, S. A., 398
Jones, C. L., 308, 1415
Jones, G. C., 173
Jones, J., 164
Jones, M. M., 1031
Jordan, R. A., 659, 2062
Joseph, S. V., 288, 868, 1918
Juan, L. W., 1087
Judd, T. M., 425

K

Kafle, L., 614
Kamble, N.A.S.a.S. T., 990
Kang, Q., 1393
Kanobe, C., 845
Karabörklü, S., 1212
Kasseney, B. D., 862
Kaufman, P. E., 250, 1108
Kaur, A., 1999
Kavallieratos, N. G., 1752, 1765
Kawahara, N., 646
Kawase, A., 646
Ke, X., 1892
Kean, J. M., 1462
Kells, S. A., 1423
Kemp, W. P., 1162
Kennedy, G. G., 143
Keremane, M. L., 586
Khamis, M., 325
Khrimian, A., 494
Kikkert, J. R., 709
Kinoshita, A., 646
Kirrane, M. J., 1146
Knight, A. L., 1309
Knodel, J. J., 1236
Ko, C.-H., 823
Koch, K. A., 1824
Koehler, G., 1613
Komatsu, J., 1969
Koppel, A. L., 137
Koul, O., 1534
Kovacs, E., 1018
Kozak, P. R., 1774
Kramer, M., 87
Krimmel, W. A., 331
Krupke, C. H., 343
Kuenen, L.p.s. B., 532
Kuhar, T. P., 137
Kullik, S. A., 484
Kumano, N., 420
Kung, Y.-L., 225
Kuniyoshi, C. H., 548
Kurashima, R., 1969
Kurihara, J., 1280
Kuriwada, T., 420
Kurtz, R., 1584

L

Lacey, I., 1999
Lacey, M. J., 232
Lafontaine, J. P., 1018
Laguna, I., 1793
Lam, A., 1093
Lambkin, T. A., 629
Lamp, W. O., 1316
Lampinen, B., 965
Lance, D. R., 494, 901
Lapointe, S. L., 540
Larson, R. A., 1633
Latham, D. R., 1864
Law, Y. H., 331
Lax, A. R., 1009
Leahy, K., 1613
Ledezma, L. A., 920, 1706
Lee, C.-Y., 232, 607, 622, 1675, 2031, 2087
Lee, D. J., 2073
Lee, R. F., 586
Lee, S. T., 1561
Lei, C., 1189
Lei, Z., 1771
Lenz, M., 232
Leonard, D. S., 1265
Leskey, T. C., 1613
Leva, N., 494
Lewis, D. S., 1986
Lewis, P., 1599
Li, A. Y., 258
Li, J., 732
Li, K., 149
Li, X., 956
Li, X.-Q., 1075
Li, Y., 1065
Li, Zhihong, 1065, 1502
Liburd, O. E., 939
Lien, Y.-Y., 225
Light, D. M., 1309
Lim, E. G., 414
Lim, U. T., 1833
Lin, C.-C., 225
Lin, H.-M., 225
Lin, Y., 1730
Ling, S., 243
Liu, B., 978
Liu, J., 1502
Liu, T.-X., 1116, 1851
Liu, X., 1384
Liu, Y., 1080
Liu, Y.-B., 1455
Liu, Z., 94
Lognay, G., 1220
Lombardo, A., 180
Long, W., 827
Lopez, R. G., 548
Los, L., 1613
Louis, J., 2068
Lu, Cheng, 101
Lu, Chong, 1189
Lu, W., 509
Lu, Z., 1730
Lucas, N., 220
Lucchi, A., 1125
Lucia, A., 595, 1087
Luna, J.-C., 1229
Lundgren, J. G., 343
Luo, C. W., 149
Luo, Jing, 1189
Luo, Junyu, 673
Luo, Y., 1892
Luthe, D. S., 299
Lye, G. C., 107
Lyons, D. B., 173

M

Ma, E., 827
Ma, Y., 673
Maccagnani, B., 266
MacGuidwin, A., 1568
MacIntosh, G. C., 845
MacKinnon, W. E., 1928
Magalhaes, L. C., 1950
Magarey, R. D., 1870
Mahroof, R. M., 1124
Mailleux, A. C., 1220
Maki, E. C., 933
Mallinger, R. E., 115
Malone, S., 137

Manandhar, R., 947
Mangan, R. L., 54, 388
Mankin, R. W., 81, 939
Mankowski, M. E., 740
Mao, L., 1002
Marco, M., 1087
Marley, K. A., 1633
Martínez, A.-M., 1229
Martini, X., 1138
Marzachì, C., 360
Marzouki, W., 188
Mashtoly, T. A., 41
Mason, C. E., 806
Massa, M. J., 1195
Mastro, V. C., 901
Masuh, H. M., 595, 1087
Mau, R.F.L., 1856
Maugin, S., 636
McCarville, M. T., 845
McCullough, D. G., 517, 1599
McInnis, D., 1969
McNeil, J. N., 94
McPheron, B. A., 1065
Medici, A., 266
Medina, P., 1541
Meeker, J. R., 1258
Meihls, L. N., 1045
Meikle, W. G., 753
Meng, J., 1189
Meyer, W. L., 939
Millar, J. G., 555, 933, 1592
Miller, D. R., 895, 1245, 1934
Mills, N. J., 1864
Miranda, D., 1316
Miranda, F., 782
Mirkov, E., 1138
Mitchell, P. D., 343
Mitchell, V. J., 1301
Mizell, R. F., 47
Mo, J., 862
Mohamed, S. A., 1180
Mohammadi, M., 1379
Monteiro, L. B., 636
Montez, G. H., 1286
Montoya, P., 1885
Moreno, A.M.T., 54, 388
Morin, G., 1613
Morrell, J. J., 740
Morse, J. G., 773
Moy, J. H., 1509
Moyseenko, J. J., 1665
Mullins, A. J., 1687
Munyaneza, J. E., 1486, 1783
Murphy, J. F., 1167
Murugan, M., 1406
Myers, S. W., 21, 220

N

Na, J. H., 693
Nabity, P. D., 459
Nakamoto, S. T., 566
Nakao, T., 646
Nam, Y., 693
Nansen, C., 1138
Naranjo, S. E., 773, 1575
Nault, B. A., 1, 1153, 1622
Navarro-Llopis, V., 1349, 1647
Nderitu, P. W., 1180
Neoh, K.-B., 607, 622
Neven, L. G., 398
Newcomb, R. D., 920
Ni, X., 956
Nielson, C., 1038
Ning, Z., 1364
Nome, C., 1793
Nomura, M., 1280
Nuessly, G. S., 155, 474
Nyrop, J. P., 1622

O

O'Brien, T. E., 1680
Odhiambo, J. F., 1584
Oghiakhe, S., 889
Okuyama, T., 2039
Olafson, P. U., 250
Olivares, J., 636
Oliver, J. B., 574, 1665, 2017
O'Neal, M., 845, 1286
Onstad, D. W., 343
Onufrieva, K., 1265
Opit, G. P., 308, 1415
O'Rourke, P. K., 1476
Ortego, F., 1349
Ortín-Angulo, M.C., 1742
Osborne, J. L., 107
Osbrink, W.L.A., 1009, 1024
Oswald, K. J., 1038
Othman, A. S., 1675
Owens, C., 1687
Oyarzún, M. P., 258

P

Pahio, E., 1969
Pakyari, H., 799
Palyvos, N. E., 1752
Pan, H., 978
Paquette, S. R., 920
Paradell, S., 1793
Park, C. G., 414
Parrella, M. P., 1323
Parsa, S., 331
Patt, J. M., 753
Paula-Moraes, S., 1900
Payton, M. E., 308
Pecci, M.P.G., 1793
Pelletier, Y., 1075
Peng, W., 732
Perera, N., 1349
Perez-Flores, L. J., 1991
Pérez-Hedo, M., 853
Perez-Mendoza, J., 443, 2078
Periera, L. N., 1093
Phillips, T. W., 1415, 1509
Picanço, M. C., 1909
Piesman, J., 659
Pineda, S., 1229
Piñero, J. C., 1613, 1856
Ping, W., 1502
Pitts-Singer, T. L., 750
Pitzer, J. B., 250, 1108
Pizzolato, T. D., 806
Pless, C. D., 665
Pluempanupat, W., 1534
Pola, S., 823
Poland, T. M., 81, 173, 517, 1599
Polat, E., 839
Pompon, J., 1075
Poopathi, S., 1816
Popay, A. J., 665
Porter, K. B., 1928
Porter, R. P., 343
Prabhaker, N., 773
Pradhan, N. C., 947
Prakash, S., 1369
Prasifka, J. R., 1561
Prezoto, F., 1120
Primo, J., 1647
Prokopy, R. J., 1613

Q

Qin, A. Z., 509

R

Raabe, M. C., 1514
Ragsdale, D. W., 1824
Raies, A., 188
Raman, A., 436
Ranger, C. M., 405, 574, 1665, 2017
Ravuiwasa, K. T., 1525
Razmjou, J., 1379
Razze, J. M., 806
Reber, J., 1055, 1393
Redak, R. A., 555
Reddy, G.V.P., 436
Reding, M. E., 405, 574, 1665, 2017
Reese, J. C., 2068
Reierson, D. A., 601
Reitz, S. R., 1771
Ren, X., 653
Rhainds, M., 1928
Riley, D. G., 956
Rinderer, T. E., 26, 1146
Rinehart, J. P., 1162
Rivera-Cabrera, F., 1991
Rizzo, R., 180
Robacker, D. C., 54, 1195
Roberts, J.M.K., 465
Robertson, J. L., 1031
Robinson, R. D., 540
Robinson, V.-A., 1229
Rodstein, J., 933
Roh, H. S., 414
Rose, C. J., 920
Rosenheim, J. A., 331

Rugman-Jones, P. F., 279
Ruiz-Arce, R., 1065
Rush, C., 1138
Rust, M. K., 601
Ryoo, M. I., 693

S

Saad, A. R., 2031
Sacchetti, P., 1195
Sadof, C. S., 548, 1960
Sagel, A., 1103
Saleem, M. A., 1343
Sanchez J.A., 1742
Sanchis, J., 1647
Sanders, W. R., 939
Sandhu, H. S., 155, 474
Sandoya, G., 299
Sauphanor, B., 636
Schaafsma, A. W., 484
Scherer, C. W., 1002
Schneider, M.-I., 1229
Schneweis, D., 1406
Scholte, E.-J., 266
Schortemeyer, M., 717
Schultz, P. B., 574, 1665, 2017
Schulze, T. L., 659, 2062
Sciarretta, r., 1752
Sears, M. K., 484
Segoli, Michal, 331
Segoli, Moran, 331
Sengoda, V. G., 1486
Serba, D. D., 2073
Shapiro-Ilan, D. I., 14, 47
Shapland, E. B., 367
Shearman, R. C., 2073
Shelly, T., 1969
Shelton, A. M., 1, 709
Shelton, C. R., 1093
Shen, J., 1364
Shen, K., 1840
Shi, G. L., 375
Shi, P., 1502
Shih, C.-J., 614
Shinde, C. K., 1369
Shiromoto, K., 420
Siegel, J. P., 532
Siegfried, B. D., 205
Siegwart, M., 636
Silk, P. J., 1928
Silva, R.S.D., 1909
Simmons, G. S., 1301
Simpson, J. A., 717
Sims, S. R., 1680
Singham, G. V., 1675
Sisterson, M. S., 367
Sivakoff, F. S., 331
Skoda, S. R., 1103
Skuhrovec, J., 1870
Smagghe, G., 1229
Smith, C. M., 1406
Soopaya, R., 1999
Soroka, J. J., 125
Souder, S. K., 1856
Souza, A.R.D., 1120
Sparks, Jr., A. N., 956
Spencer, J. L., 343
Spurgeon, D. W., 685
Stamm, M. D., 205
Stanley, C. A., 750
Starkey, S., 1406
Steinmann, K. P., 1496
Stelinski, L. L., 540, 939
Stephen, P. R., 1174, 1180
Stephens, A.E.A., 1999
Stevens, B. R., 1986
Stouthamer, R., 279
Strange, J. P., 732
Straub, R. W., 1622
Stringer, L. D., 1301, 1462, 1999
Strom, B. L., 1258
Stuart, J. J., 1055
Su, Jianya, 653, 1364
Su, Jun, 1592
Su, N.-Y., 1687
Su, Q., 978
Su, Y.-T., 823
Subramanyam, B., 317, 325
Suckling, D. M., 920, 1301, 1462, 1514, 1999
Sukhirun, N., 1534
Sullivan, N. J., 1301
Sullivan, T.E.S., 1301
Sun, J.-R., 1080
Sun, Y. Y., 149
Sutherland, A. M., 1323
Suthisut, D., 1445
Sutton, C., 220
Sweeney, J. D., 1928

T

Tan, C.-W., 1525
Tantravahi, P., 1138
Tasin, M., 1125
Taverner, P. D., 220
Teal, P.E.A., 1430
Teale, S. A., 1592
Tee, H.-S., 2031
Tenbroeck, S. H., 1108
Thomas, D. B., 54, 63, 388
Thomas, K., 717
Throne, J. E., 443, 1765, 2078
Tian, M. Y., 509
Tobin, P. C., 1265
Tooker, J. F., 343
Tooman, L. K., 920
Toscano, N. C., 773
Toyoshima, G., 1280
Trematerra, P., 1752
Troxclair, N., 1138
Trumble, J. T., 586
Tsai, C.-W., 1480
Tsai, W.-T., 274
Tseng, Y.-C., 225
Tsunoda, K., 232
Tuna, G., 839
Tuttle, A., 1613

U

Ugine, T. A., 1942
Ulyshen, M. D., 81, 501
Umina, P. A., 465
Uzunovic, A., 717

V

Vacas, S., 1647
Vandervoort, C., 1656
Vankosky, M. A., 1550
Vargas, R. I., 1856
Varón, E., 164
Vassiliou, V. A., 580, 1979
Vaughn, K., 1138
Vazquez-Luna, A., 1991
Vega, F. E., 87
Venâncio, D.F.A., 1120
Verde, G. L., 180
Vetter, R. S., 601, 986
Vijayaraghavan, R., 1369
Villa, J. D., 745
Viñuela, E., 1229, 1541
Virla, E., 1793
Vorsino, A., 947
Vreysen, M.j.b., 814

W

Wagnitz, J., 1146
Wagstaff, S. J., 1514
Walgenbach, J. F., 1950
Walton, V. M., 970
Wang, B., 1720, 1730
Wang, C., 274
Wang, D., 1720
Wang, J.-J., 1436, 1720
Wang, K.-H., 947
Wang, P., 709
Wang, Q., 509
Wang, S., 978
Wang, X.Z.H., 1640
Wang, Y. N., 375
Wang, Z.-Y., 1720
Waterworth, R. A., 555
Wearing, C. H., 1514
Webb, S. E., 155, 474
Weeks, A. R., 465
Wei, D.-D., 1720
Weinzierl, R. A., 1031
Werf, W.V.D., 673
Werling, B. P., 1622
West, M., 1162
Whelan, P. M., 1146
White, J., 274
Whitfield, A. E., 1406
Wickham, J. D., 1592
Williams, P., 299
Willis, D. K., 1800
Wilson, L. T., 702
Wise, J. C., 1656

Wistrom, C. M., 367
Witzgall, P., 1125
Wong, N., 2087
Wood, B. W., 14
Woods, B., 1462, 1999
Workneh, F., 1138
Wright, M. G., 947
Wright, R. J., 1900
Wu, B., 1840
Wu, D., 833
Wu, Jiajiao, 1065
Wu, Jie, 732
Wu, K., 94, 833
Wu, Qingjun, 978
Wu, Quancong, 1877
Wu, S., 2046
Wu, W.-J., 2039
Wu, Y., 1065
Wyckhuys, K.A.G., 833

X

Xia, J., 673
Xiao, P., 1877
Xiao, W., 1592
Xie, W., 978
Xu, B., 978
Xu, H., 1730
Xu, J., 509
Xu, Z.-H., 1080
Xue, Y., 1138

Y

Yan, R., 1892
Yang, N., 978
Yang, X., 978
Yang, Y., 702, 1730
Ye, G., 1892, 2046
Yee, W. L., 194, 909
Yilmaz, S., 1212
Yocum, G. D., 1162
Yokoyama, V. Y., 792
Yoshimura, T., 232
Youssef, N., 1665
Yu, C., 317
Yu, Q.-Y., 101
Yuan, M.-L., 1720
Yue, B., 1116

Z

Zalom, F., 965
Zangerl, A. R., 459
Zanuncio, J. C., 1120
Zaviezo, T., 331
Zee, F. T., 566, 571
Zerba, E. N., 595, 1087
Zewdu, A., 1329
Zhang, A., 1265
Zhang, F., 1592
Zhang, H., 1055
Zhang, J., 1877
Zhang, M., 1496
Zhang, R., 243, 1840
Zhang, Yongmei, 1116
Zhang, Youjun, 978
Zhang, Ze, 101
Zhang, Zhilin, 1189
Zhao, B., 1189
Zhao, J., 833
Zhao, K., 1502
Zhao, X., 1364
Zheng, Xianyun, 827
Zheng, Xuesong, 653
Zheng, Xusong, 1730
Zhou, D., 1502
Zhu, L., 1384, 2068
Zhu, Y. C., 1364, 1877
Zuo, W.-D., 101

CUMULATIVE KEY WORD INDEX
Journal of Economic Entomology
Volume 104, 2011

A

above and belowground interactions,
aphid colonization, 1568
Aphid glycines, 1568
Heterodera glycines, 1568
path analysis, 1568
Acalymma trivittatum,
bait, 1294
chemical control, 1294
Diabrotica undecimpunctata undecimpunctata, 1294
Acari,
Acaridae, 1423
disinfestation, 1423
nonchemical control, 1423
supercooling point, 1423
acaricidal activity,
Inula japonica, 375
Tetranychus cinnabarinus, 375
Acaridae,
Acari, 1423
disinfestation, 1423
nonchemical control, 1423
supercooling point, 1423
accession,
aphid, 1075
breeding, 1075
potato, 1075
resistance, 1075
accuracy,
efficiency, 1323
Frankliniella occidentalis, 1323
precision, 1323
sampling, 1323
acetic acid,
grape juice, 1195
methyl anthranilate, 1195
Mexican fruit fly, 1195
propylene glycol, 1195
Adelges tsugae,
distribution, 1918
eastern hemlock, 1918
tree height, 1918
Tsuga canadensis, 1918
Adelphocoris suturalis,
circadian rhythm, 1189
Y-tube bioassay, 1189
adjuvant,
insecticide management, 1309
kairomone, 1309
organophosphate, 1309
pyrethroid, 1309
Adress,
Ceratitis capitata, 1647
chemosterilant, 1647
lufenuron, 1647
SIT, 1647
Aedes aegypti,
amide, 1369
behavioral response, 1369
mosquito, 1369
repellent, 1369
aeration,
insects, 702
modeling, 702
rice, 702
storage, 702
Aethina tumida, 26, 753
diet, 753
in-hive traps, 26
out-hive traps, 26
oviposition, 753
polyvinyl chloride trap, 26
survivorship, 753
temperature, 753
white trap, 26
age-stage,
Chilades pandava peripatria, 1525
Cycas revoluta, 1525
Cycas taitungensis, 1525
two-sex life table, 1525
aggregation,
integrated pest management, 1613
monitoring, 1613
reduced-risk, 1613
semiochemicals, 1613
aggregation pheromone,
2-(undecyloxy)-ethanol, 1592
ethanol, 1592
Monochamus alternatus, 1592
α-pinene, 1592
agriculture,
Bombus terrestris, 107
pollination limitation, 107
Rubus idaeus, 107
Scotland, 107
Agrilus,
Fraxinus, 69
fumigation, 69
methyl bromide, 69
quarantine, 69
Agrilus planipennis, 81, 379
auditory cues, 81
biocontrol, 379
cryptic, 81
parasitoid rearing, 379
Spathius agrili, 379
vibrokinesis, 81
vibrotaxis, 81
agroforestry systems,
Coffea arabica, 164
leaf-cutting ants, 164
vegetational diversity, 164
albedo,
flavedo, 54
host status, 54
Rutaceae, 54
Tephritidae, 54
alcohol,
cockroach, 1680
oil, 1680
synergism, 1680
toxicity, 1680
Aleurothrixus floccosus,
citrus, 1329
cyhalothrin, 1329
selecron, 1329
woolly whiteflies, 1329
alkaloid,
corn flea beetle, 665
endophyte, 665
insect resistance, 665
Neotyphodium, 665
Allee effect,
biological invasion, 1928
competitive displacement, 1928
Allium cepa, 1, 1622
cultural control, 1622
Delia antiqua, 1622
Iris yellow spot virus, 1
management, 1
oviposition, 1622
planting date, 1622
Thrips tabaci, 1
allyl isothiocyanate,
ambrosia beetles, 1665
botanical insecticides, 1665
essential oils, 1665
Xylosandrus germanus, 1665
almond leaf scorch,
almonds, 367
Draeculacephala minerva, 367
sharpshooter, 367
vector management, 367
almonds, 211, 367, 965
almond leaf scorch, 367
Amyelois transitella, 211, 965
Draeculacephala minerva, 367
hull split, 965
monitoring, 211
navel orangeworm, 965
oviposition traps, 211
pistachios, 211
sharpshooter, 367
shell seal, 965
vector management, 367
Alphitobius diaperinus,
cross-resistance, 629
darkling beetle, 629
metabolic resistance, 629
piperonyl butoxide, 629

Alpinia conchigera,
contact toxicity, 1445
Curcuma zedoaria, 1445
synthetic oil, 1445
Zingiber zerumbet, 1445
Alpinia galanga,
Bactrocera dorsalis, 1534
carboxylesterase, 1534
glutathione transferase, 1534
toxicity, 1534
alternative pollinator,
cold storage, 1162
fluctuating thermal regime, 1162
fluctuation thermal regime, 1162
Megachile rotundata, 1162
Amblyomma americanum, 659, 2062
carvacrol, 659
EcoSMART, 659
Ixodes scapularis, 659, 2062
nootkatone, 659
repellent testing, 2062
ambrosia beetles, 574, 1665
allyl isothiocyanate, 1665
botanical insecticides, 1665
essential oils, 1665
ethanol, 574
Scolytinae, 574
Xylosandrus germanus, 1665
α-pinene, 574
American elm,
bioassay, 889
Dutch elm disease, 889
efficacy, 889
native elm bark beetle, 889
American serpentine leafminer,
ionizing radiation, 1851
phytosanitary treatment, 1851
quarantine treatment, 1851
amide,
Aedes aegypti, 1369
behavioral response, 1369
mosquito, 1369
repellent, 1369
Ampelovirus,
Closteroviridae, 1480
Planococcus ficus, 1480
semipersistent, 1480
Amyelois transitella, 211, 965
almonds, 211, 965
hull split, 965
monitoring, 211
navel orangeworm, 965
oviposition traps, 211
pistachios, 211
shell seal, 965
Anastrepha,
competitive displacement, 1204
host status, 1204
infestation patterns, 1204
Psidium guajava, 1204
Anastrepha fruit flies,
Citrus, 388
commodity quarantines, 388
fruit fly hosts, 388
Anastrepha ludens, 63, 452
commodity treatment, 63
fruit fly, 452
malathion, 452
phytosanitary, 63
quarantine, 63
spinosad, 452
traps, 452
Anoplophora glabripennis,
fecundity, 1942
invasive species, 1942
neonicotinoid, 1942
sublethal effect, 1942
antibiosis,
coevolution, 1055
effector-triggered immunity, 1055
gall midges, 1055
anticoagulants,
Cochliomyia hominivorax, 1103
defibrination 1103
insect diet, 1103
antioxidant defenses,
Cd^{2+}, 827
lipid peroxidation, 827
oxidative stress, 827
Propsilocerus akamusi, 827
antitermitic activity,
Bacillus subtilis subsp. *virginiana*, 1502
biocontrol, 1502
termite, 1502
ants,
aphid, 947
Banana bunchy top virus, 947
sampling, 947
within-plant distribution, 947
aphid, 947, 1075, 1167
accession, 1075
ants, 947
Banana bunchy top virus, 947
barley yellow dwarf, 1167
breeding, 1075
cereal yellow dwarf, 1167
potato, 1075
resistance, 1075
Rhopalosiphum padi, 1167
sampling, 947
virus vectors, 1167
within-plant distribution, 947
aphid biotype,
Sitobion avenae, 1080
wheat, 1080
wheat resistance to aphid, 1080
aphid colonization,
above and belowground interactions, 1568
Aphid glycines, 1568
Heterodera glycines, 1568
path analysis, 1568
Aphid glycines,
above and belowground interactions, 1568
aphid colonization, 1568
Heterodera glycines, 1568
path analysis, 1568
aphid-bioassay,
Aphis glycines, 1357
insecticide susceptibility, 1357
soybean aphid, 1357
Aphis glycines, 1357, 1824
aphid-bioassay, 1357
Entomophthorales, 1824
Glycine max, 1824
insecticide susceptibility, 1357
nontarget impacts, 1824
Rag1, 1824
soybean aphid, 1357
Aphis gossypii,
Cucumis sativus, 1379
life table, 1379
plant resistance, 1379
Apis mellifera,
breeding, 745
mite resistance, 745
Varroa destructor, 745
apterae,
extraction, 1116
poplar petiolegall aphid, 1116
sampling, 1116
spatial distribution, 1116
Arachnida,
Holocnemus pluchei, 601
pest control, 601
urban entomology, 601
artificial rearing,
bark beetles, 87
host-marking pheromones, 87
Hypothenemus hampei, 87
ash canopy,
green prism traps, 173
monitoring, 173
phoebe oil, 173
Z-3-hexenol, 173
ash tree protection,
dinotefuran, 1599
emerald ash borer, 1599
imidacloprid, 1599
invasive pest, 1599
aster leafhopper,
aster yellows, 1800
aster yellows phytoplasma, 1800
Macrosteles quadrilineatus, 1800
qPCR, 1800
aster yellows,
aster leafhopper, 1800
aster yellows phytoplasma, 1800
Macrosteles quadrilineatus, 1800
qPCR, 1800
aster yellows phytoplasma,
aster leafhopper, 1800
aster yellows, 1800
Macrosteles quadrilineatus, 1800
qPCR, 1800
asynchronous development,
mating failure, 1146
nonreproduction, 1146
resistance, 1146
Varroa destructor, 1146
attractant,
Cerambycidae, 1258
monitoring, 1258

trap design, 1258
woodborer, 1258
auditory cues,
Agrilus planipennis, 81
cryptic, 81
vibrokinesis, 81
vibrotaxis, 81
avirulence gene,
coevolution, 1393
Mayetiola destructor, 1393
plant breeding, 1393
pyramiding, 1393

B

B. thuringiensis subsp. *Israelensis*,
Bacillus sphaericus, 1816
biopesticides, 1816
Coffee husk waste, 1816
mosquitoes, 1816
Bacillus sphaericus,
B. thuringiensis subsp. *Israelensis*, 1816
biopesticides, 1816
Coffee husk waste, 1816
mosquitoes, 1816
Bacillus subtilis subsp. *virginiana*,
antitermitic activity, 1502
biocontrol, 1502
termite, 1502
Bacillus thuringiensis, 673, 1045, 1476
cotton, 673
cowpea trypsin inhibitor, 673
Diabrotica virgifera virgifera, 1045
Helicoverpa armigera, 673
Helicoverpa zea, 1476
MIR604, 1045
Ostrinia nubilalis, 1476
resistance, 673
resistance management, 1476
toxicity assay, 1045
Bacillus thuringiensis kurstaki and *aizawai*,
black cutworm, 41
Noctuidae, 41
bacterial challenge,
glutathione S-transferase, 2046
homology modeling, 2046
insecticides, 2046
mRNA expression, 2046
Bactrocera cucurbitae, 566, 571, 1065, 1509
Bactrocera dorsalis, 566, 571
Bactrocera latifrons, 571, 1509
Ceratitis capitata, 566, 571
gene flow, 1065
genetic structure, 1065
microsatellites, 1065
phytosanitary treatment, 1509
quarantine pest, 1065, 1509
resistance, 566
Vaccinium spp., 566, 571
X-ray irradiation, 1509
Bactrocera dorsalis, 566, 571, 1534, 1969
Alpinia galanga, 1534
Bactrocera cucurbitae, 566, 571
Bactrocera latifrons, 571
carboxylesterase, 1534
Ceratitis capitata, 566, 571
glutathione transferase, 1534
invasive species, 1969
mating competitiveness, 1969
resistance, 566
sterile male releases, 1969
toxicity, 1534
Vaccinium spp., 566, 571
Bactrocera invadens,
cold disinfestation, 1180
commodity treatment, 1180
phytosanitary, 1180
quarantine, 1180
Bactrocera latifrons, 571, 1509
Bactrocera cucurbitae, 571, 1509
Bactrocera dorsalis, 571
Ceratitis capitata, 571
phytosanitary treatment, 1509
quarantine pest, 1509
Vaccinium spp., 571
X-ray irradiation, 1509
Bactrocera tau,
development, 1840
host species, 1840
intraspecific competition, 1840
larval density, 1840
bait,
Acalymma trivittatum, 1294
chemical control, 1294
Diabrotica undecimpunctata undecimpunctata, 1294
bait station,
behavior, 1856
integrated pest management, 1856
Mediterranean fruit fly, 1856
rainfastness, 1856
bait trap station,
detection dog, 225
fire ant detection, 225
odor recognition, 225
red imported fire ant, 225
baiting,
bistrifluron, 607
colony elimination, 607
Globitermes sulphureus, 607
secondary pest, 607
Banana bunchy top virus,
ants, 947
aphid, 947
sampling, 947
within-plant distribution, 947
barcodes,
internal transcribed spacer, 1706
real-time polymerase chain reaction, 1706
Tortricidae, 1706
bark beetles,
artificial rearing, 87
host-marking pheromones, 87
Hypothenemus hampei, 87
barley yellow dwarf,
aphids, 1167
cereal yellow dwarf, 1167
Rhopalosiphum padi, 1167
virus vectors, 1167
base temperature,
day degrees, 1870
lower developmental threshold, 1870
rate of development, 1870
sum of effective temperatures, 1870
baseline susceptibility, 653, 1950
bioassay, 653
Cnaphalocrocis medinalis, 653
codling moth, 1950
insecticide resistance, 653
oriental fruit moth, 1950
reduced-risk insecticides, 1950
Beauveria bassiana,
carbaryl, 14
Curculio caryae, 14
cypermethrin, 14
Steinernema, 14
bed bug,
Cimex lectularius, 274
dry ice, 274
monitoring, 274
traps, 274
bee dispersal,
bee retention, 750
Osmia lignaria, 750
solitary bees, 750
bee retention,
bee dispersal, 750
Osmia lignaria, 750
solitary bees, 750
behavior,
bait station, 1856
integrated pest management, 1856
Mediterranean fruit fly, 1856
rainfastness, 1856
behavioral response,
Aedes aegypti, 1369
amide, 1369
mosquito, 1369
repellent, 1369
Bemisia tabaci, 548, 833, 978
biotype B, 978
biotype Q, 978
China, 978
displacement, 978
Euphorbia pulcherrima, 548
imidacloprid, 833
poinsettia, 548
sublethal effect, 833
sweetpotato whitefly, 548
toxicity, 833
beneficial insects,
egg parasitoids, 773
laboratory bioassay, 773

predators, 773
systemic insecticides, 773
benefit-cost ratio,
binomial sampling, 1900
resampling software, 1900
western bean cutworm, 1900
binomial sampling,
benefit-cost ratio, 1900
resampling software, 1900
western bean cutworm, 1900
bioassay, 653, 889, 1031, 2095
American elm, 889
baseline susceptibility, 653
Cnaphalocrocis medinalis, 653
Dutch elm disease, 889
efficacy, 889
Grapholita molesta, 1031
insecticide resistance, 653
Ixodes scapularis, 2095
mealworms, 2095
Metarhizium brunneum, 2095
native elm bark beetle, 889
resistance monitoring, 1031
Tenebrio molitor, 2095
bioassays residual activity,
chronic exposure, 405
neonicotinoids, 405
Otiorhynchus sulcatus, 405
biochemical mechanism,
bisultap resistant strain, 243
cross-resistance, 243
Nilaparvata lugens, 243
biocontrol, 379, 1502
Agrilus planipennis, 379
antitermitic activity, 1502
Bacillus subtilis subsp. *virginiana*, 1502
parasitoid rearing, 379
Spathius agrili, 379
termite, 1502
bioenergy,
defoliator, 1561
ethanol, 1561
pest, 1561
resistance, 1561
biofuels,
fall armyworm, 459
herbivory, 459
leaf toughness, 459
biological control, 32, 94, 343, 693, 799, 1496, 1742, 2031
Bt corn, 343
Chrysoperla sinica, 94
cockroach control, 2031
dispersal, 94
fecundity, 799
flight activity, 94
flight mill, 94
host age preference, 2031
insect resistance management, 343
integrated pest management, 1496, 1742
intraspecific competition, 693
intrinsic rate of increase, 799
leafcutter ants, 32
life cycle, 32
mirid, 1742
monitoring, 1742
native phorid species, 32
oothecal parasitoid, 2031
parasitism rate, 32
pear psyllid, 1742
Periplaneta americana, 2031
Plodia interpunctella, 693
population growth, 799
refuge, 343
secondary pest outbreaks, 1496
secondary pests, 343
Sitophilus oryzae, 693
temperature threshold, 799
Tetranychidae, 1496
tree nuts, 1496
biological invasion, 1265, 1771, 1928
Allee effect, 1928
competitive displacement, 1928
insecticide susceptibility, 1771
IPM, 1265
Liriomyza sativae, 1771
Liriomyza trifolii, 1771
Lymantria dispar, 1265
pheromone release rate, 1265
pheromone-baited traps, 1265
species displacement, 1771
biomarker,
glutathione transferase, 101
oxidative stress, 101
phoxim, 101
silkworm, 101
biomolecules,
fruit fly, 1991
graft, 1991
mango, 1991
preharvest, 1991
biopesticides,
B. thuringiensis subsp. *Israelensis*, 1816
Bacillus sphaericus, 1816
Coffee husk waste, 1816
mosquitoes, 1816
biorational,
larvicide, 1986
methionine, 1986
biotype,
Diuraphis noxia, 1736
host plant resistance, 1736
Russian wheat aphid, 1736
biotype B,
Bemisia tabaci, 978
biotype Q, 978
China, 978
displacement, 978
biotype Q,
Bemisia tabaci, 978
biotype B, 978
China, 978
displacement, 978
bistrifluron,
baiting, 607
colony elimination, 607
Globitermes sulphureus, 607
secondary pest, 607
bisultap resistant strain,
biochemical mechanism, 243
cross-resistance, 243
Nilaparvata lugens, 243
black cutworm,
Bacillus thuringiensis kurstaki and *aizawai*, 41
Noctuidae, 41
Blattella germanica,
essential oils, 595
Eucalyptus, 595
knockdown, 595
monoterpenes, 595
Blattisocius tarsalis,
Cryptolestes ferrugineus, 1752
Lepidoglyphus destructor, 1752
Plodia interpunctella, 1752
SADIE, 1752
blended seed refuge,
European corn borer, 806
feeding behavior, 806
resistance management, 806
transgenic corn, 806
Blissus occiduus, 205, 2073
Buchloe dactyloides, 2073
buffalograss, 205
chinch bug, 205
LC_{50}, 205
neonicotinoids, 205
plant resistance, 2073
turfgrass, 2073
Bombus terrestris,
agriculture, 107
pollination limitation, 107
Rubus idaeus, 107
Scotland, 107
borneol,
1-octen-3-ol, 1633
menthol, 1633
repellency, 1633
volatiles, 1633
botanical insecticides, 1665, 1979
allyl isothiocyanate, 1665
ambrosia beetles, 1665
damage, 1979
essential oils, 1665
grapefruit, 1979
Kelly's citrus thrips, 1979
organic, 1979
Pezothrips kellyanus, 1979
Xylosandrus germanus, 1665
bottle-traps,
ethanol concentration, 2017
Xylosandrus crassiusculus, 2017
Xylosandrus germanus, 2017
Brassicae oleracea,
Diadegma insulare, 782
Nicaragua, 782
Plutella xylostella, 782
Polybia sp., 782
breeding, 745, 1075
accession, 1075
aphid, 1075
Apis mellifera, 745
mite resistance, 745
potato, 1075

resistance, 1075
Varroa destructor, 745
broad bean,
chrysanthemum yellows, 360
Flavescence dorée, 360
leafhopper, 360
phytoplasma transmission, 360
Bt,
Diabrotica virgifera virgifera, 1584
eCry3.1Ab, 1584
mCry3A, 1584
resistance management, 1584
Bt corn, 343, 853
biological control, 343
development, 853
ecdysteroids, 853
insect resistance management, 343
juvenile hormone, 853
refuge, 343
secondary pests, 343
Sesamia nonagrioides, 853
Bt resistance,
Diabrotica virgifera virgifera, 1038
insect resistance management, 1038
MON863, 1038
Buchloe dactyloides,
Blissus occiduus, 2073
plant resistance, 2073
turfgrass, 2073
buffalograss,
Blissus occiduus, 205
chinch bug, 205
LC_{50}, 205
neonicotinoids, 205
bumble bee,
honey bee, 732
host expansion, 732
virus, 732
bunch damage,
Cyprus, 580
insecticides, 580
Lobesia botrana, 580
vineyard, 580
Buprestidae,
Cerambycidae, 1934
Curculionidae, 1934
Monochamus titillator, 1934
Scolytinae, 1934
burnt cane,
green cane, 155
lesser cornstalk borer, 155
sugarcane, 155
tillage, 155
butene-fipronil,
cross-resistance, 1364
ethiprole, 1364
fipronil, 1364
Nilaparvata lugens, 1364

C

C. modoc,
C. vicinus, 740
carpenter ants, 740
colony founding, 740
relative humidity, 740
C. vicinus,
C. modoc, 740
carpenter ants, 740
colony founding, 740
relative humidity, 740
Calepitrimerus vitis,
integrated pest management, 970
predatory mites, 970
sublethal effects, 970
toxicity, 970
canola,
flea beetle feeding, 125
host plant resistance, 125
Phyllotreta spp., 125
trichomes, 125
carbaryl,
Beauveria bassiana, 14
Curculio caryae, 14
cypermethrin, 14
Steinernema, 14
carbohydrates,
lipids, 1229
methoxyfenozide, 1229
proteins, 1229
Spodoptera exigua, 1229
carbon dioxide,
environment, 1774
modified atmosphere, 1774
mortality, 1774
Varroa destructor, 1774
carboxylesterase,
Alpinia galanga, 1534
Bactrocera dorsalis, 1534
glutathione transferase, 1534
toxicity, 1534
carpenter ants,
C. modoc, 740
C. vicinus, 740
colony founding, 740
relative humidity, 740
carvacrol,
Amblyomma americanum, 659
EcoSMART, 659
Ixodes scapularis, 659
nootkatone, 659
cations,
foraging, 425
Reticulitermes flavipes, 425
soil, 425
termite, 425
causal inference,
ecoinformatics, 331
economic injury level, 331
observational studies, 331
statistical power, 331
Cd^{2+},
antioxidant defenses, 827
lipid peroxidation, 827
oxidative stress, 827
Propsilocerus akamusi, 827
Cerambycidae, 1258, 1934
attractant, 1258
Buprestidae, 1934
Curculionidae, 1934
monitoring, 1258
Monochamus titillator, 1934
Scolytinae, 1934
trap design, 1258
woodborer, 1258
Ceratitis capitata, 566, 571, 1174, 1647
Adress, 1647
Bactrocera cucurbitae, 566, 571
Bactrocera dorsalis, 566, 571
Bactrocera latifrons, 571
chemosterilant, 1647
cold disinfestation, 1174
commodity treatment, 1174
lufenuron, 1647
phytosanitary regulation, 1174
quarantine, 1174
resistance, 566
SIT, 1647
Vaccinium spp., 566, 571
cereal yellow dwarf,
aphids, 1167
barley yellow dwarf, 1167
Rhopalosiphum padi, 1167
virus vectors, 1167
chalkbrood,
foulbrood, 353
honey bees, 353
ozone, 353
pathogens, 353
chemical control, 137, 1294, 1349, 1885
Acalymma trivittatum, 1294
bait, 1294
Diabrotica undecimpunctata undecimpunctata, 1294
egg efficacy, 137
esterases, 1349
fruit fly programs, 1885
integrated pest management, 1885
Mediterranean fruit fly, 1349
Pentatomidae, 137
resistance management, 1349
Scelionidae, 137
synergist, 1349
chemical cues,
Coleoptera, 436
Cosmopolites sordidus, 436
Curculionidae, 436
visual cues, 436
chemosterilant,
Adress, 1647
Ceratitis capitata, 1647
lufenuron, 1647
SIT, 1647
Chilades pandava peripatria,
age-stage, 1525
Cycas revoluta, 1525

Cycas taitungensis, 1525
two-sex life table, 1525
China,
Bemisia tabaci, 978
biotype B, 978
biotype Q, 978
displacement, 978
chinch bug,
Blissus occiduus, 205
buffalograss, 205
LC_{50}, 205
neonicotinoids, 205
chlorantraniliprole,
indoxacarb, 990
persistence, 990
R. flavipes, 990
termiticide, 990
chlorfluazuron,
diflubenzuron, 1009
eastern subterranean termite, 1009
Formosan termite, 1009
hexaflumuron, 1009
chronic exposure,
bioassays residual activity, 405
neonicotinoids, 405
Otiorhynchus sulcatus, 405
chrysanthemum yellows,
broad bean, 360
Flavescence dorée, 360
leafhopper, 360
phytoplasma transmission, 360
chrysanthemyl 2-methylbutanoate,
field bioassay, 823
lavandulyl 2-methylbutanoate, 823
Madeira mealybug, 823
stereochemistry, 823
Chrysoperla sinica,
biological control, 94
dispersal, 94
flight activity, 94
flight mill, 94
Cimex lectularius,
bed bug, 274
dry ice, 274
monitoring, 274
traps, 274
circadian rhythm,
Adelphocoris suturalis, 1189
Y-tube bioassay, 1189
citral,
common thyme, 414
essential oils, 414
lemongrass, 414
quarantine, 414
citrus,
Aleurothrixus floccosus, 1329
cyhalothrin, 1329
selecron, 1329
woolly whiteflies, 1329
Citrus,
Anastrepha fruit flies, 388
commodity quarantines, 388
fruit fly hosts, 388
citrus leafminer,
sex pheromone, 540
SPLAT, 540
(Z,Z)-7,11-hexadecadienal, 540
(Z,Z,*E*)-7,11,13-hexadecatrienal, 540
citrus peelminer,
degree-day, 1286
lower developmental threshold, 1286
Closteroviridae,
Ampelovirus, 1480
Planococcus ficus, 1480
semipersistent, 1480
clothianidin,
compensatory overconsumption, 484
hormesis, 484
hormoligosis, 484
toxin interaction, 484
Cnaphalocrocis medinalis, 653, 1730
baseline susceptibility, 653
bioassay, 653
Cry1C*, 1730
Cry2A*, 1730
insecticide resistance, 653
resistance, 1730
rice, 1730
Cochliomyia hominivorax,
anticoagulants, 1103
defibrination, 1103
insect diet, 1103
cockroach,
alcohol, 1680
oil, 1680
synergism, 1680
toxicity, 1680
cockroach control,
biological control, 2031
host age preference, 2031
oothecal parasitoid, 2031
Periplaneta americana, 2031
codling moth, 814, 1950
baseline susceptibility, 1950
long-distance transportation, 814
oriental fruit moth, 1950
reduced-risk insecticides, 1950
sterile insect technique, 814
coevolution, 1055, 1393
antibiosis, 1055
avirulence gene, 1393
effector-triggered immunity, 1055
gall midges, 1055
Mayetiola destructor, 1393
plant breeding, 1393
pyramiding, 1393
Coffea arabica,
agroforestry systems, 164
leaf-cutting ants, 164
vegetational diversity, 164
coffee,
coffee insect pests, 1909
conventional cultivation, 1909
organic cultivation, 1909
semiochemical, 1909
Coffee husk waste,
B. thuringiensis subsp. *Israelensis*, 1816
Bacillus sphaericus, 1816
biopesticides, 1816
mosquitoes, 1816
coffee insect pests,
coffee, 1909
conventional cultivation, 1909
organic cultivation, 1909
semiochemical, 1909
cold disinfestation, 1174, 1180
Bactrocera invadens, 1180
Ceratitis capitata, 1174
commodity treatment, 1174, 1180
phytosanitary, 1180
phytosanitary regulation, 1174
quarantine, 1174, 1180
cold storage, 1162, 1833
alternative pollinator, 1162
conservational biological control, 1833
fluctuating thermal regime, 1162
fluctuation thermal regime, 1162
Gryon japonicum, 1833
Megachile rotundata, 1162
Ooencyrtus nezarae, 1833
stink bug, 1833
cold treatment,
heat treatment, 21
hot water immersion, 21
phytosanitary treatment, 21
quarantine treatment, 21
Coleoptera,
chemical cues, 436
Cosmopolites sordidus, 436
Curculionidae, 436
visual cues, 436
colony elimination,
baiting, 607
bistrifluron, 607
Globitermes sulphureus, 607
secondary pest, 607
colony founding,
C. modoc, 740
C. vicinus, 740
carpenter ants, 740
relative humidity, 740
commodity quarantines,
Anastrepha fruit flies, 388
Citrus, 388
fruit fly hosts, 388
commodity treatment, 63, 1174, 1180
Anastrepha ludens, 63
Bactrocera invadens, 1180
Ceratitis capitata, 1174
cold disinfestation, 1174, 1180
phytosanitary, 63, 1180
phytosanitary regulation, 1174
quarantine, 63, 1174, 1180
common eastern bumble bee,
European honey bee, 1153
foraging behavior, 1153

P. pruinosa, 1153
pumpkin, 1153
common smut,
corn earworm, 956
edible oils, 956
husk coverage, 956
sap beetles, 956
common thyme,
citral, 414
essential oils, 414
lemongrass, 414
quarantine, 414
compensation,
CP 78-1628, 474
CP 88-1762, 474
CP 89-2143, 474
lesser cornstalk borer, 474
compensatory overconsumption,
clothianidin, 484
hormesis, 484
hormoligosis, 484
toxin interaction, 484
competitive attraction,
mating disruption, 939
pheromone dispenser, 939
Vitacea polistiformis, 939
competitive displacement, 1204, 1928
Allee effect, 1928
Anastrepha, 1204
biological invasion, 1928
host status, 1204
infestation patterns, 1204
Psidium guajava, 1204
conservational biological control,
cold storage, 1833
Gryon japonicum, 1833
Ooencyrtus nezarae, 1833
stink bug, 1833
contact toxicity,
Alpinia conchigera, 1445
Curcuma zedoaria, 1445
synthetic oil, 1445
Zingiber zerumbet, 1445
Contarinia nasturtii,
integrated pest management, 709
invasive species, 709
North America, 709
control, 839, 1087
Eucalyptus, 1087
Haematobia irritans, 1087
horn flies, 1087
insect growth regulator, 839
knockdown, 1087
Lycoriella ingenua, 839
mushroom sciarid fly, 839
soil drench, 839
controlled atmosphere temperature treatment system,
heat, 398
Macchiademus diplopterus, 398
Phlyctinus callosus, 398
postharvest, 398
conventional cultivation,
coffee, 1909
coffee insect pests, 1909
organic cultivation, 1909
semiochemical, 1909
Coptotermes formosanus, 232, 1002, 2025
delayed action, 2025
lethal contraction, 1002
lethal dose, 1002
nonrepellent, 2025
paper consumption, 232
Reticulitermes flavipes, 1002
Reticulitermes speratus, 232
substrate type, 2025
termite lipid content, 232
termite survival, 232
termiticide, 2025
toxicity ranking, 1002
corn earworm,
common smut, 956
edible oils, 956
husk coverage, 956
sap beetles, 956
corn flea beetle,
alkaloid, 665
endophyte, 665
insect resistance, 665
Neotyphodium, 665
corn leafhopper,
corn stunt, 1793
Dalbulus maidis, 1793
spiroplasmas, 1793
corn stunt,
corn leafhopper, 1793
Dalbulus maidis, 1793
spiroplasmas, 1793
Cosmopolites sordidus,
chemical cues, 436
Coleoptera, 436
Curculionidae, 436
visual cues, 436
cotton, 673, 685
Bacillus thuringiensis, 673
cowpea trypsin inhibitor, 673
Helicoverpa armigera, 673
plant bug, 685
resistance, 673
sampling, 685
sweep net, 685
cowpea trypsin inhibitor,
Bacillus thuringiensis, 673
cotton, 673
Helicoverpa armigera, 673
resistance, 673
CP 78-1628,
compensation, 474
CP 88-1762, 474
CP 89-2143, 474
lesser cornstalk borer, 474
CP 88-1762,
compensation, 474
CP 78-1628, 474
CP 89-2143, 474
lesser cornstalk borer, 474
CP 89-2143,
compensation, 474
CP 78-1628, 474
CP 88-1762, 474
lesser cornstalk borer, 474
crop damage,
Lygus elisus, 1575
Lygus lineolaris, 1575
Lygus hesperus, 1575
Physaria fendleri, 1575
cross-resistance, 243, 629, 1364
Alphitobius diaperinus, 629
biochemical mechanism, 243
bisultap resistant strain, 243
butene-fipronil, 1364
darkling beetle, 629
ethiprole, 1364
fipronil, 1364
metabolic resistance, 629
Nilaparvata lugens, 243, 1364
piperonyl butoxide, 629
Cry1C*,
Cnaphalocrocis medinalis, 1730
Cry1C*, 1730
Cry2A*, 1730
resistance, 1730
rice, 1730
Cry2A*,
Cnaphalocrocis medinalis, 1730
Cry1C*, 1730
Cry2A*, 1730
resistance, 1730
rice, 1730
cryptic,
Agrilus planipennis, 81
auditory cues, 81
vibrokinesis, 81
vibrotaxis, 81
Cryptolestes ferrugineus,
Blattisocius tarsalis, 1752
Lepidoglyphus destructor, 1752
Plodia interpunctella, 1752
SADIE, 1752
Cryptotermes,
de-alates, 622
nymphoid, 622
sex determination, 622
sex ratio, 622
Cucumis sativus,
Aphis gossypii, 1379
life table, 1379
plant resistance, 1379
cultural control,
Allium cepa, 1622
Delia antiqua, 1622
oviposition, 1622
planting date, 1622
Curculio caryae,
Beauveria bassiana, 14
carbaryl, 14
cypermethrin, 14
Steinernema, 14
Curculionidae, 436, 1934
Buprestidae, 1934
Cerambycidae, 1934
chemical cues, 436

Coleoptera, 436
Cosmopolites sordidus, 436
Monochamus titillator, 1934
Scolytinae, 1934
visual cues, 436
Curcuma zedoaria,
Alpinia conchigera, 1445
contact toxicity, 1445
synthetic oil, 1445
Zingiber zerumbet, 1445
Cycas revoluta,
age-stage, 1525
Chilades pandava peripatria, 1525
Cycas taitungensis, 1525
two-sex life table, 1525
Cycas taitungensis,
age-stage, 1525
Chilades pandava peripatria, 1525
Cycas revoluta, 1525
two-sex life table, 1525
Cydia molesta,
insecticide resistance, 636
mechanism, 636
monitoring, 636
target mutation, 636
cyhalothrin,
Aleurothrixus floccosus, 1329
citrus, 1329
selecron, 1329
woolly whiteflies, 1329
cypermethrin, 14, 258
Beauveria bassiana, 14
carbaryl, 14
Curculio caryae, 14
diazinon, 258
horn fly, 258
insecticide resistance, 258
kdr, 258
Steinernema, 14
Cyprus,
bunch damage, 580
insecticides, 580
Lobesia botrana, 580
vineyard, 580
cytochrome C oxidase I,
28S, 279
Graphocephala cythura, 279
Graphocephala flavovittata, 279
sharpshooters, 279

D

dairy,
house fly, 1093
IPM, 1093
monitoring, 1093
surveillance, 1093
Dalbulus maidis,
corn leafhopper, 1793
corn stunt, 1793
spiroplasmas, 1793
damage, 180, 1979
botanical insecticides, 1979
gall density, 180
grapefruit, 1979
kaolin, 180
Kelly's citrus thrips, 1979
Ophelimus maskelli, 180
organic, 1979
Pezothrips kellyanus, 1979
darkling beetle,
Alphitobius diaperinus, 629
cross-resistance, 629
metabolic resistance, 629
piperonyl butoxide, 629
Dasineura mali,
parasitism, 1640
phenology, 1640
Platygaster demades, 1640
synchronization, 1640
day degrees,
base temperature, 1870
lower developmental threshold, 1870
rate of development, 1870
sum of effective temperatures, 1870
de-alates,
Cryptotermes, 622
nymphoid, 622
sex determination, 622
sex ratio, 622
defibrination,
anticoagulants, 1103
Cochliomyia hominivorax, 1103
insect diet, 1103
defoliator,
bioenergy, 1561
ethanol, 1561
pest, 1561
resistance, 1561
degree-day, 532, 1286
citrus peelminer, 1286
development, 532
lower developmental threshold, 1286
navel orangeworm, 532
pistachios, 532
survival, 532
delayed action,
Coptotermes formosanus, 2025
nonrepellent, 2025
substrate type, 2025
termiticide, 2025
delayed differential equation model,
Epiphyas postvittana, 1462
eradication, 1462
irradiation biology, 1462
Delia antiqua,
Allium cepa, 1622
cultural control, 1622
oviposition, 1622
planting date, 1622
detection,
mealybugs, 555
ornamental plants, 555
pheromone trap, 555
sampling, 555
detection dog,
bait trap station, 225
fire ant detection, 225
odor recognition, 225
red imported fire ant, 225
detection tools,
double-decker trap, 517
green leaf volatiles, 517
invasive forest pest, 517
manuka oil, 517
development, 532, 853, 1436, 1840
Bactrocera tau, 1840
Bt corn, 853
degree-days, 532
ecdysteroids, 853
host species, 1840
intraspecific competition, 1840
juvenile hormone, 853
larval density, 1840
life table, 1436
navel orangeworm, 532
pistachios, 532
psocid, 1436
Sesamia nonagrioides, 853
stored products, 1436
survival, 532
temperature, 1436
development rate,
fecundity, 1864
population growth rate, 1864
thermal requirements, 1864
developmental period,
flour, 2087
population growth, 2087
starch, 2087
Tribolium castaneum, 2087
Deverra scoparia,
essential oil, 1220
fecundity, 1220
Tetranychus urticae, 1220
toxicity, 1220
Diabrotica undecimpunctata undecimpunctata,
Acalymma trivittatum, 1294
bait, 1294
chemical control, 1294
Diabrotica virgifera virgifera, 1038, 1045, 1584, 2054
Bacillus thuringiensis, 1045
Bt, 1584
Bt resistance, 1038
eCry3.1Ab, 1584
insect resistance management, 1038, 2054
mCry3A, 1584, 2054
MIR604, 1045
MON863, 1038
phenology, 2054
resistance management, 1584
toxicity assay, 1045
Diadegma insulare,
Brassicae oleracea, 782
Nicaragua, 782
Plutella xylostella, 782
Polybia sp., 782

diazinon,
cypermethrin, 258
horn fly, 258
insecticide resistance, 258
kdr, 258
diet,
Aethina tumida, 753
oviposition, 753
survivorship, 753
temperature, 753
diflubenzuron,
chlorfluazuron, 1009
eastern subterranean termite, 1009
Formosan termite, 1009
hexaflumuron, 1009
3,5-dimethyldodecanoic acid,
pest management, 933
pheromone, 933
dinotefuran,
ash tree protection, 1599
emerald ash borer, 1599
imidacloprid, 1599
invasive pest, 1599
disinfection,
eggs, 188
Mediterranean fruit fly, 188
quaternary ammonium, 188
rearing, 188
disinfestation,
Acari, 1423
Acaridae, 1423
nonchemical control, 1423
supercooling point, 1423
dispersal,
biological control, 94
Chrysoperla sinica, 94
flight activity, 94
flight mill, 94
displacement,
Bemisia tabaci, 978
biotype B, 978
biotype Q, 978
China, 978
distribution,
Adelges tsugae, 1918
eastern hemlock, 1918
tree height, 1918
Tsuga canadensis, 1918
disturbance,
edge, 879
exotic species, 879
survey, 879
trapping, 879
Diuraphis noxia,
biotype, 1736
host plant resistance, 1736
Russian wheat aphid, 1736
diurnal pollination,
female reproductive success, 149
Jatropha curcas, 149
nectar production, 149
nocturnal pollination, 149
diversity,
Eucalyptus, 1120
Polistinae, 1120
DNA barcoding,
internal transcribed spacer, 1720
Psocoptera, 1720
species identification, 1720
double-decker trap,
detection tools, 517
green leaf volatiles, 517
invasive forest pest, 517
manuka oil, 517
Draeculacephala minerva,
almond leaf scorch, 367
almonds, 367
sharpshooter, 367
vector management, 367
Drosophila,
natural lure, 1018
trap, 1018
dry ice,
bed bug, 274
Cimex lectularius, 274
monitoring, 274
traps, 274
Dutch elm disease,
American elm, 889
bioassay, 889
efficacy, 889
native elm bark beetle, 889

E

eastern hemlock, 868, 1918
Adelges tsugae, 1918
distribution, 1918
fertilizer, 868
hemlock woolly adelgid, 868
imidacloprid, 868
predators, 868
tree height, 1918
Tsuga canadensis, 1918
eastern subterranean termite,
chlorfluazuron, 1009
diflubenzuron, 1009
Formosan termite, 1009
hexaflumuron, 1009
ecdysteroids,
Bt corn, 853
development, 853
juvenile hormone, 853
Sesamia nonagrioides, 853
ecoinformatics,
causal inference, 331
economic injury level, 331
observational studies, 331
statistical power, 331
economic analysis,
insecticide coverage, 1960
Oligonycus aceris, 1960
secondary pest outbreak, 1960
Xylosandrus crassiusculus, 1960
economic injury level,
causal inference, 331
ecoinformatics, 331
observational studies, 331
statistical power, 331
economic threshold,
field pea, 1550
nitrogen fertilizer, 1550
pea leaf weevil, 1550
yield loss, 1550
EcoSMART,
Amblyomma americanum, 659
carvacrol, 659
Ixodes scapularis, 659
nootkatone, 659
eCry3.1Ab,
Bt, 1584
Diabrotica virgifera virgifera, 1584
mCry3A, 1584
resistance management, 1584
edge,
disturbance, 879
exotic species, 879
survey, 879
trapping, 879
edible oils,
common smut, 956
corn earworm, 956
husk coverage, 956
sap beetles, 956
effector-triggered immunity,
antibiosis, 1055
coevolution, 1055
gall midges, 1055
efficacy,
American elm, 889
bioassay, 889
Dutch elm disease, 889
native elm bark beetle, 889
efficiency,
accuracy, 1323
Frankliniella occidentalis, 1323
precision, 1323
sampling, 1323
egg efficacy,
chemical control, 137
Pentatomidae, 137
Scelionidae, 137
egg parasitoids,
beneficial insects, 773
laboratory bioassay, 773
predators, 773
systemic insecticides, 773
eggs,
disinfection, 188
Mediterranean fruit fly, 188
quaternary ammonium, 188
rearing, 188
elytral color polymorphism,
mass-marking method, 420
mating compatibility, 420
sterile insect technique, 420
emerald ash borer, 901, 1599
ash tree protection, 1599
dinotefuran, 1599
funnel trap, 901
imidacloprid, 1599
invasive pest, 1599
prism trap, 901

trap color, 901
visual cues, 901
emergence,
insect pest, 465
life cycle, 465
Sminthurus viridis, 465
soil moisture, 465
Empoasca fabae,
host plant expansion, 1316
photosynthesis, 1316
transpiration, 1316
enantiomers,
Ips avulsus, 895
ipsdienol, 895
seasonal variation, 895
semiochemical context, 895
endophyte,
alkaloid, 665
corn flea beetle, 665
insect resistance, 665
Neotyphodium, 665
Entomophthorales,
Aphis glycines, 1824
Glycine max, 1824
nontarget impacts, 1824
Ragl, 1824
environment,
carbon dioxide, 1774
modified atmosphere, 1774
mortality, 1774
Varroa destructor, 1774
EPG,
feeding behavior, 2068
resistance, 2068
soybean aphids, 2068
Ephestia kuehniella,
fecundity, 1212
fumigant toxicity, 1212
longevity, 1212
Epiphyas postvittana, 220, 920, 1301, 1462, 1999
delayed differential equation model, 1462
eradication, 1301, 1462, 1999
haplotype, 920
inherited sterility, 1999
invasive species, 920
irradiation, 1999
irradiation biology, 1462
mitochondrial population genetics, 920
origin, 920
ovicidal activity, 220
petroleum oils, 220
radiation, 1301
reduced risk pesticides, 220
sterile insect technique, 1301, 1999
Tortricidae, 1301
eradication, 1301, 1462, 1999
delayed differential equation model, 1462
Epiphyas postvittana, 1301, 1462, 1999
inherited sterility, 1999
irradiation, 1999
irradiation biology, 1462
radiation, 1301
sterile insect technique, 1301, 1999
Tortricidae, 1301
essential oil, 414, 595, 1220, 1665
allyl isothiocyanate, 1665
ambrosia beetles, 1665
Blattella germanica, 595
botanical insecticides, 1665
citral, 414
common thyme, 414
Deverra scoparia, 1220
Eucalyptus, 595
fecundity, 1220
knockdown, 595
lemongrass, 414
monoterpenes, 595
quarantine, 414
Tetranychus urticae, 1220
toxicity, 1220
Xylosandrus germanus, 1665
esterases,
chemical control, 1349
Mediterranean fruit fly, 1349
resistance management, 1349
synergist, 1349
ethanol, 574, 1245, 1561, 1592
2-(undecyloxy)-ethanol, 1592
aggregation pheromone, 1592
ambrosia beetles, 574
bioenergy, 1561
defoliator, 1561
ipsdienol, 1245
ipsenol, 1245
Monochamus alternatus, 1592
Monochamus titillator, 1245
pest, 1561
α-pinene, 574, 1245, 1592
resistance, 1561
Scolytinae, 574
ethanol concentration,
bottle-traps, 2017
Xylosandrus crassiusculus, 2017
Xylosandrus germanus, 2017
ethiprole,
butene-fipronil, 1364
cross-resistance, 1364
fipronil, 1364
Nilaparvata lugens, 1364
Eucalyptus, 595, 1087, 1120
Blattella germanica, 595
control, 1087
diversity, 1120
essential oils, 595
Haematobia irritans, 1087
horn flies, 1087
knockdown, 595, 1087
monoterpenes, 595
Polistinae, 1120
Euphorbia pulcherrima,
Bemisia tabaci, 548
poinsettia, 548
sweetpotato whitefly, 548
European corn borer,
blended seed refuge, 806
feeding behavior, 806
resistance management, 806
transgenic corn, 806
European honey bee,
common eastern bumble bee, 1153
foraging behavior, 1153
P. pruinosa, 1153
pumpkin, 1153
exotic,
invasive, 501
Lauraceae, 501
laurel wilt, 501
sassafras, 501
exotic species,
disturbance, 879
edge, 879
survey, 879
trapping, 879
extraction,
apterae, 1116
poplar petiolegall aphid, 1116
sampling, 1116
spatial distribution, 1116

F

fall armyworm, 299, 459
biofuels, 459
herbivory, 459
leaf toughness, 459
Lepidoptera, 299
maize, 299
plant–herbivore interactions, 299
rootworm, 299
fecundity, 509, 799, 1212, 1220, 1864, 1942
Anoplophora glabripennis, 1942
biological control, 799
development rate, 1864
Deverra scoparia, 1220
Ephestia kuehniella, 1212
essential oil, 1220
fumigant toxicity, 1212
intrinsic rate of increase, 799
invasive species, 1942
longevity, 509, 1212
neonicotinoid, 1942
phenology, 509
population growth, 799
population growth rate, 1864
rearing, 509
sublethal effect, 1942
survival, 509
temperature threshold, 799
Tetranychus urticae, 1220
thermal requirements, 1864
toxicity, 1220
feeding behavior, 806, 2068
blended seed refuge, 806
EPG, 2068
European corn borer, 806
resistance, 2068
resistance management, 806

soybean aphids, 2068
transgenic corn, 806
feeding rates,
monitoring station, 1024
seasonal trends, 1024
subterranean termite, 1024
female reproductive success,
diurnal pollination, 149
Jatropha curcas, 149
nectar production, 149
nocturnal pollination, 149
fertility,
hemlock woolly adelgid, 288
T. canadensis, 288
T. heterophylla, 288
T. mertensiana, 288
fertilizer,
eastern hemlock, 868
hemlock woolly adelgid, 868
imidacloprid, 868
predators, 868
field bioassay,
chrysanthemyl 2-methylbutanoate, 823
lavandulyl 2-methylbutanoate, 823
Madeira mealybug, 823
stereochemistry, 823
field pea,
economic threshold, 1550
nitrogen fertilizer, 1550
pea leaf weevil, 1550
yield loss, 1550
fipronil, 1272, 1364, 1541
butene-fipronil, 1364
cross-resistance, 1364
ethiprole, 1364
imidacloprid, 1272, 1541
Nantucket pine tip moth, 1272
Nilaparvata lugens, 1364
pales weevil, 1272
Psyttalia concolor, 1541
side effects, 1541
systemic insecticides, 1272
fipronil resistance,
GABA receptor, 646
Laodelphax striatellus, 646
fire ant detection,
bait trap station, 225
detection dog, 225
odor recognition, 225
red imported fire ant, 225
flavedo,
albedo, 54
host status, 54
Rutaceae, 54
Tephritidae, 54
Flavescence dorée,
broad bean, 360
chrysanthemum yellows, 360
leafhopper, 360
phytoplasma transmission, 360
flea beetle feeding,
canola, 125
host plant resistance, 125
Phyllotreta spp., 125
trichomes, 125
flight activity,
biological control, 94
Chrysoperla sinica, 94
dispersal, 94
flight mill, 94
flight initiation, 443, 2078
food deprivation, 443
mating status, 2078
presence or absence of food, 2078
rearing density, 443
red flour beetle, 443
starvation, 2078
Tribolium castaneum, 443, 2078
flight mill,
biological control, 94
Chrysoperla sinica, 94
dispersal, 94
flight activity, 94
flour,
developmental period, 2087
population growth, 2087
starch, 2087
Tribolium castaneum, 2087
fluctuating thermal regime,
alternative pollinator, 1162
cold storage, 1162
fluctuation thermal regime, 1162
Megachile rotundata, 1162
fluctuation thermal regime,
alternative pollinator, 1162
cold storage, 1162
fluctuating thermal regime, 1162
Megachile rotundata, 1162
Folsomia candida,
nontargets, 1892
risk assessment, 1892
transgenic Bt rice, 1892
food deprivation,
flight initiation, 443
rearing density, 443
red flour beetle, 443
Tribolium castaneum, 443
foraging,
cations, 425
Reticulitermes flavipes, 425
soil, 425
termite, 425
foraging behavior,
common eastern bumble bee, 1153
European honey bee, 1153
P. pruinosa, 1153
pumpkin, 1153
forest defoliator,
invasive species, 494
pheromone trap survey, 494
Formosan termite,
chlorfluazuron, 1009
diflubenzuron, 1009
eastern subterranean termite, 1009
hexaflumuron, 1009
foulbrood,
chalkbrood, 353
honey bees, 353
ozone, 353
pathogens, 353
Frankliniella occidentalis,
accuracy, 1323
efficiency, 1323
precision, 1323
sampling, 1323
Fraxinus,
Agrilus, 69
fumigation, 69
methyl bromide, 69
quarantine, 69
fruit fly, 452, 1991
Anastrepha ludens, 452
biomolecules, 1991
graft, 1991
malathion, 452
mango, 1991
preharvest, 1991
spinosad, 452
traps, 452
fruit fly hosts,
Anastrepha fruit flies, 388
Citrus, 388
commodity quarantines, 388
fruit fly programs,
chemical control, 1885
integrated pest management, 1885
fumigant toxicity,
Ephestia kuehniella, 1212
fecundity, 1212
longevity, 1212
fumigation, 69, 792
Agrilus, 69
Fraxinus, 69
hydrogen phosphide, 792
Mayetiola destructor, 792
methyl bromide, 69
Oulema melanopus, 792
pressure, 792
quarantine, 69
funnel trap,
emerald ash borer, 901
prism trap, 901
trap color, 901
visual cues, 901

G

GABA receptor,
fipronil resistance, 646
Laodelphax striatellus, 646
gall density,
damage, 180
kaolin, 180
Ophelimus maskelli, 180
gall midges,
antibiosis, 1055
coevolution, 1055
effector-triggered immunity, 1055

gene flow,
Bactrocera cucurbitae, 1065
genetic structure, 1065
microsatellites, 1065
quarantine pest, 1065
generating susceptible strain,
insecticide resistance, 1343
resistance management, 1343
Spodoptera exigua, 1343
genetic structure,
Bactrocera cucurbitae, 1065
gene flow, 1065
microsatellites, 1065
quarantine pest, 1065
Globitermes sulphureus,
baiting, 607
bistrifluron, 607
colony elimination, 607
secondary pest, 607
glutathione S-transferase,
bacterial challenge, 2046
homology modeling, 2046
insecticides, 2046
mRNA expression, 2046
glutathione transferase, 101, 1534
Alpinia galanga, 1534
Bactrocera dorsalis, 1534
biomarker, 101
carboxylesterase, 1534
oxidative stress, 101
phoxim, 101
silkworm, 101
toxicity, 1534
Glycine max,
Aphis glycines, 1824
Entomophthorales, 1824
nontarget impacts, 1824
Rag1, 1824
graft,
biomolecules, 1991
fruit fly, 1991
mango, 1991
preharvest, 1991
grape juice,
acetic acid, 1195
methyl anthranilate, 1195
Mexican fruit fly, 1195
propylene glycol, 1195
grapefruit,
botanical insecticides, 1979
damage, 1979
Kelly's citrus thrips, 1979
organic, 1979
Pezothrips kellyanus, 1979
grapes,
Japanese beetle, 1656
rainfastness, 1656
residual activity, 1656
residue profile, 1656
Graphocephala cythura,
28S, 279
cytochrome C oxidase I, 279
Graphocephala flavovittata, 279
sharpshooters, 279
Graphocephala flavovittata,
28S, 279
cytochrome C oxidase I, 279
Graphocephala cythura, 279
sharpshooters, 279
Grapholita molesta,
bioassays, 1031
resistance monitoring, 1031
green cane,
burnt cane, 155
lesser cornstalk borer, 155
sugarcane, 155
tillage, 155
green leaf volatiles,
detection tools, 517
double-decker trap, 517
invasive forest pest, 517
manuka oil, 517
green prism traps,
ash canopy, 173
monitoring, 173
phoebe oil, 173
Z-3-hexenol, 173
Gryon japonicum,
cold storage, 1833
conservational biological control, 1833
Ooencyrtus nezarae, 1833
stink bug, 1833

H

Haematobia irritans,
control, 1087
Eucalyptus, 1087
horn flies, 1087
knockdown, 1087
haplotype,
Epiphyas postvittana, 920
invasive species, 920
mitochondrial population genetics, 920
origin, 920
heat,
controlled atmosphere temperature treatment system, 398
Macchiademus diplopterus, 398
Phlyctinus callosus, 398
postharvest, 398
heat tolerance,
heat treatment, 317
methyl bromide alternative, 317
heat treatment, 21, 317, 1415
cold treatment, 21
heat tolerance, 317
hot water immersion, 21
Liposcelis corrodens, 1415
Liposcelis decolor, 1415
methyl bromide alternative, 317
phytosanitary treatment, 21
quarantine treatment, 21
Rhyzopertha dominica, 1415
Tribolium castaneum, 1415
Helicoverpa armigera,
Bacillus thuringiensis, 673
cotton, 673
cowpea trypsin inhibitor, 673
resistance, 673
Helicoverpa zea,
Bacillus thuringiensis, 1476
Ostrinia nubilalis, 1476
resistance management, 1476
hemlock woolly adelgid, 288, 868
eastern hemlock, 868
fertility, 288
fertilizer, 868
imidacloprid, 868
predators, 868
T. canadensis, 288
T. heterophylla, 288
T. mertensiana, 288
herbivore-induced plant volatile,
lacewing, 115
methyl salicylate, 115
soybean aphid, 115
syrphid fly, 115
herbivory,
biofuels, 459
fall armyworm, 459
leaf toughness, 459
Hessian fly,
Mayetiola destructor, 1384
phytohormones, 1384
rice, 1384
wheat, 1384
Heterodera glycines,
above and belowground interactions, 1568
aphid colonization, 1568
Aphid glycines, 1568
path analysis, 1568
Heterorhabditis,
peach, 47
Prunus persicae, 47
Steinernema carpocapsae, 47
Synanthedon pictipes, 47
Heterotermes aureus,
monitors, 1339
subterranean termites, 1339
trapping, 1339
hexaflumuron,
chlorfluazuron, 1009
diflubenzuron, 1009
eastern subterranean termite, 1009
Formosan termite, 1009
Holocnemus pluchei,
Arachnida, 601
pest control, 601
urban entomology, 601
homology modeling,
bacterial challenge, 2046
glutathione S-transferase, 2046
insecticides, 2046
mRNA expression, 2046
honey bee, 353, 732
bumble bee, 732
chalkbrood, 353
foulbrood, 353
host expansion, 732
ozone, 353

pathogens, 353
virus, 732
hormesis,
clothianidin, 484
compensatory overconsumption, 484
hormoligosis, 484
toxin interaction, 484
hormoligosis,
clothianidin, 484
compensatory overconsumption, 484
hormesis, 484
toxin interaction, 484
horn fly, 258, 1087
control, 1087
cypermethrin, 258
diazinon, 258
Eucalyptus, 1087
Haematobia irritans, 1087
insecticide resistance, 258
kdr, 258
knockdown, 1087
host age preference,
biological control, 2031
cockroach control, 2031
oothecal parasitoid, 2031
Periplaneta americana, 2031
host expansion,
bumble bee, 732
honey bee, 732
virus, 732
host plant expansion,
Empoasca fabae, 1316
photosynthesis, 1316
transpiration, 1316
host plant resistance, 125, 143, 1736
biotype, 1736
canola, 125
Diuraphis noxia, 1736
flea beetle feeding, 125
Phyllotreta spp., 125
Russian wheat aphid, 1736
sweetpotato, 143
sweetpotato flea beetle, 143
trichomes, 125
wireworm, 143
host range,
host specificity, 1514
invasive species, 1514
pest risk analysis, 1514
Tortricidae, 1514
host species,
Bactrocera tau, 1840
development, 1840
intraspecific competition, 1840
larval density, 1840
host specificity,
host range, 1514
invasive species, 1514
pest risk analysis, 1514
Tortricidae, 1514
host status, 54, 1204
albedo, 54
Anastrepha, 1204
competitive displacement, 1204
flavedo, 54
infestation patterns, 1204
Psidium guajava, 1204
Rutaceae, 54
Tephritidae, 54
host-marking pheromones,
artificial rearing, 87
bark beetles, 87
Hypothenemus hampei, 87
hot melt pressure sensitive adhesive,
sticky gel, 909
trap design, 909
western cherry fruit fly, 909
yellow color, 909
hot water immersion,
cold treatment, 21
heat treatment, 21
phytosanitary treatment, 21
quarantine treatment, 21
house fly,
dairy, 1093
IPM, 1093
monitoring, 1093
surveillance, 1093
hull split,
almonds, 965
Amyelois transitella, 965
navel orangeworm, 965
shell seal, 965
husk coverage,
common smut, 956
corn earworm, 956
edible oils, 956
sap beetles, 956
hydrogen phosphide,
fumigation, 792
Mayetiola destructor, 792
Oulema melanopus, 792
pressure, 792
Hypothenemus hampei,
artificial rearing, 87
bark beetles, 87
host-marking pheromones, 87

I

imidacloprid, 833, 868, 1272, 1541, 1599
ash tree protection, 1599
Bemisia tabaci, 833
dinotefuran, 1599
eastern hemlock, 868
emerald ash borer, 1599
fertilizer, 868
fipronil, 1272, 1541
hemlock woolly adelgid, 868
invasive pest, 1599
Nantucket pine tip moth, 1272
pales weevil, 1272
predators, 868
Psyttalia concolor, 1541
side effects, 1541
sublethal effect, 833
systemic insecticides, 1272
toxicity, 833
indoxacarb,
chlorantraniliprole, 990
persistence, 990
R. flavipes, 990
termiticide, 990
infestation levels,
light intensity, 1675
mound size, 1675
parasitism frequencies, 1675
infestation patterns,
Anastrepha, 1204
competitive displacement, 1204
host status, 1204
Psidium guajava, 1204
inherited sterility,
Epiphyas postvittana, 1999
eradication, 1999
irradiation, 1999
sterile insect technique, 1999
in-hive traps,
Aethina tumida, 26
out-hive traps, 26
polyvinyl chloride trap, 26
white trap, 26
inoculation efficiency,
liberibacter, 1486
potato, 1486
potato psyllid, 1486
zebra chip, 1486
insect control,
ozone, 308
psocids, 308
stored grain beetles, 308
stored wheat, 308
insect diet,
anticoagulants, 1103
Cochliomyia hominivorax, 1103
defibrination, 1103
insect growth regulator, 839, 1765
control, 839
Liposcelis, 1765
Lycoriella ingenua, 839
mushroom sciarid fly, 839
Psocoptera, 1765
pyriproxyfen, 1765
soil drench, 839
surface treatment, 1765
insect pest,
emergence, 465
life cycle, 465
Sminthurus viridis, 465
soil moisture, 465
insect resistance,
alkaloid, 665
corn flea beetle, 665
endophyte, 665
Neotyphodium, 665
insect resistance management, 343, 1038, 2054
biological control, 343
Bt corn, 343
Bt resistance, 1038
Diabrotica virgifera virgifera, 1038, 2054
mCry3A, 2054
MON863, 1038

phenology, 2054
refuge, 343
secondary pests, 343
insect vision,
LED, 1430
pitfall trapping, 1430
stored product, 1430
insecticide coverage,
economic analysis, 1960
Oligonycus aceris, 1960
secondary pest outbreak, 1960
Xylosandrus crassiusculus, 1960
insecticide management,
adjuvant, 1309
kairomone, 1309
organophosphate, 1309
pyrethroid, 1309
insecticide resistance, 258, 636, 653, 1343, 2039
baseline susceptibility, 653
bioassay, 653
Cnaphalocrocis medinalis, 653
Cydia molesta, 636
cypermethrin, 258
diazinon, 258
generating susceptible strain, 1343
horn fly, 258
kdr, 258
life history traits, 2039
mechanism, 636
monitoring, 636
naled, 2039
oriental fruit fly, 2039
population dynamics, 2039
resistance management, 1343
Spodoptera exigua, 1343
target mutation, 636
insecticide spraying,
integrated pest management, 1138
potato psyllids, 1138
row crops, 1138
water-sensitive spray cards, 1138
insecticide susceptibility, 1357, 1771
aphid-bioassay, 1357
Aphis glycines, 1357
biological invasion, 1771
Liriomyza sativae, 1771
Liriomyza trifolii, 1771
soybean aphid, 1357
species displacement, 1771
insecticides, 580, 586, 2046
bacterial challenge, 2046
bunch damage, 580
Cyprus, 580
glutathione S-transferase, 2046
homology modeling, 2046
IPM, 586
Lobesia botrana, 580
mRNA expression, 2046
potato, 586
potato psyllid, 586
vineyard, 580
zebra chip, 586
insects,
aeration, 702
modeling, 702
rice, 702
storage, 702
integrated pest management, 325, 709, 970, 1125, 1138, 1496, 1613, 1687, 1742, 1856, 1885
aggregation, 1613
bait station, 1856
behavior, 1856
biological control, 1496, 1742
Calepitrimerus vitis, 970
chemical control, 1885
Contarinia nasturtii, 709
fruit fly programs, 1885
insecticide spraying, 1138
invasive species, 709, 1687
Mediterranean fruit fly, 1856
mirid, 1742
monitoring, 1613, 1742
natural enemies, 1125
nonchemical control, 325
North America, 709
pear psyllid, 1742
pheromone mating disruption, 1125
plant volatiles, 1125
potato psyllids, 1138
predatory mites, 970
rainfastness, 1856
reduced-risk, 1613
row crops, 1138
secondary pest outbreaks, 1496
semiochemicals, 1613
stored grain, 325
sublethal effects, 970
subterranean termite, 1687
succession ecology, 1687
Tetranychidae, 1496
toxicity, 970
tree nuts, 1496
water-sensitive spray cards, 1138
interaction,
resistance, 1406
wheat, 1406
wheat curl mite, 1406
wheat streak mosaic virus, 1406
internal transcribed spacer, 1706, 1720
barcodes, 1706
DNA barcoding, 1720
Psocoptera, 1720
real-time polymerase chain reaction, 1706
species identification, 1720
Tortricidae, 1706
International Plant Protection Convention,
International Standard for Phytosanitary Measures No. 15, 717
quarantine treatments, 717
International Standard for Phytosanitary Measures No. 15,
International Plant Protection Convention, 717
quarantine treatments, 717
interspecific competition,
Monomorium chinense, 614
Pheidole fervens, 614
Solenopsis invicta, 614
Solenopsis invicta virus-1, 614
intraspecific competition, 693, 1840
Bactrocera tau, 1840
biological control, 693
development, 1840
host species, 1840
larval density, 1840
Plodia interpunctella, 693
Sitophilus oryzae, 693
intrinsic rate of increase,
biological control, 799
fecundity, 799
population growth, 799
temperature threshold, 799
Inula japonica,
acaricidal activity, 375
Tetranychus cinnabarinus, 375
invasive,
exotic, 501
Lauraceae, 501
laurel wilt, 501
sassafras, 501
invasive forest pest,
detection tools, 517
double-decker trap, 517
green leaf volatiles, 517
manuka oil, 517
invasive insect species,
Kalman filter, 764
pests, 764
state-space model, 764
invasive pest,
ash tree protection, 1599
dinotefuran, 1599
emerald ash borer, 1599
imidacloprid, 1599
invasive species, 494, 709, 920, 1514, 1687, 1942, 1969
Anoplophora glabripennis, 1942
Bactrocera dorsalis, 1969
Contarinia nasturtii, 709
Epiphyas postvittana, 920
fecundity, 1942
forest defoliator, 494
haplotype, 920
host range, 1514
host specificity, 1514
integrated pest management, 709, 1687
mating competitiveness, 1969
mitochondrial population genetics, 920
neonicotinoid, 1942
North America, 709
origin, 920
pest risk analysis, 1514

pheromone trap survey, 494
sterile male releases, 1969
sublethal effect, 1942
subterranean termite, 1687
succession ecology, 1687
Tortricidae, 1514
ionizing radiation,
American serpentine leafminer, 1851
phytosanitary treatment, 1851
quarantine treatment, 1851
IPM, 586, 845, 1093, 1265
biological invasions, 1265
dairy, 1093
house fly, 1093
insecticides, 586
Lymantria dispar, 1265
monitoring, 1093
natural enemy, 845
pheromone release rate, 1265
pheromone-baited traps, 1265
population dynamics, 845
potato, 586
potato psyllid, 586
surveillance, 1093
zebra chip, 586
Ips avulsus,
enantiomers, 895
ipsdienol, 895
seasonal variation, 895
semiochemical context, 895
ipsdienol, 895, 1245
enantiomers, 895
ethanol, 1245
Ips avulsus, 895
ipsenol, 1245
Monochamus titillator, 1245
α-pinene, 1245
seasonal variation, 895
semiochemical context, 895
ipsenol,
ethanol, 1245
ipsdienol, 1245
Monochamus titillator, 1245
α-pinene, 1245
Iris yellow spot virus,
Allium cepa, 1
management, 1
Thrips tabaci, 1
irradiation,
Epiphyas postvittana, 1999
eradication, 1999
inherited sterility, 1999
sterile insect technique, 1999
irradiation biology,
delayed differential equation model, 1462
Epiphyas postvittana, 1462
eradication, 1462
Ixodes scapularis, 659, 2062, 2095
Amblyomma americanum, 659, 2062
bioassay, 2095
carvacrol, 659
EcoSMART, 659
mealworms, 2095
Metarhizium brunneum, 2095
nootkatone, 659
repellent testing, 2062
Tenebrio molitor, 2095

J

Japanese beetle,
grapes, 1656
rainfastness, 1656
residual activity, 1656
residue profile, 1656
Jatropha curcas,
diurnal pollination, 149
female reproductive success, 149
nectar production, 149
nocturnal pollination, 149
juvenile hormone,
Bt corn, 853
development, 853
ecdysteroids, 853
Sesamia nonagrioides, 853

K

kairomone,
adjuvant, 1309
insecticide management, 1309
organophosphate, 1309
pyrethroid, 1309
Kalman filter,
invasive insect species, 764
pests, 764
state-space model, 764
kaolin,
damage, 180
gall density, 180
Ophelimus maskelli, 180
kdr,
cypermethrin, 258
diazinon, 258
horn fly, 258
insecticide resistance, 258
Kelly's citrus thrips,
botanical insecticides, 1979
damage, 1979
grapefruit, 1979
Kelly's citrus thrips, 1979
organic, 1979
Pezothrips kellyanus, 1979
Klason lignin,
Odontotermes formosanus, 862
termite feeding, 862
wood extractive, 862
wood hardness, 862
knockdown, 595, 1087
Blattella germanica, 595
control, 1087
essential oils, 595
Eucalyptus, 595, 1087
Haematobia irritans, 1087
horn flies, 1087
monoterpenes, 595

L

laboratory bioassay,
beneficial insects, 773
egg parasitoids, 773
predators, 773
systemic insecticides, 773
lacewing,
herbivore-induced plant volatile, 115
methyl salicylate, 115
soybean aphid, 115
syrphid fly, 115
Laodelphax striatellus, 646, 1877
fipronil resistance, 646
GABA receptor, 646
Nephotettix cincticeps, 1877
Nilaparvata lugens, 1877
pymetrozine, 1877
Sogatella furcifera, 1877
larval density,
Bactrocera tau, 1840
development, 1840
host species, 1840
intraspecific competition, 1840
larval development,
mosquito control, 266
pupation rate, 266
sterile insect technique, 266
survival rate, 266
larvicide,
biorational, 1986
methionine, 1986
Lauraceae,
exotic, 501
invasive, 501
laurel wilt, 501
sassafras, 501
laurel wilt,
exotic, 501
invasive, 501
Lauraceae, 501
sassafras, 501
lavandulyl 2-methylbutanoate,
chrysanthemyl 2-methylbutanoate, 823
field bioassay, 823
Madeira mealybug, 823
stereochemistry, 823
LC_{50},
Blissus occiduus, 205
buffalograss, 205
chinch bug, 205
neonicotinoids, 205
leaf toughness,
biofuels, 459
fall armyworm, 459
herbivory, 459
leafcutter ants,
biological control, 32
life cycle, 32
native phorid species, 32
parasitism rate, 32
leaf-cutting ants,
agroforestry systems, 164
Coffea arabica, 164
vegetational diversity, 164

leafhopper,
broad bean, 360
chrysanthemum yellows, 360
Flavescence dorée, 360
phytoplasma transmission, 360
LED,
insect vision, 1430
pitfall trapping, 1430
stored product, 1430
lemongrass,
citral, 414
common thyme, 414
essential oils, 414
quarantine, 414
Lepidoglyphus destructor,
Blattisocius tarsalis, 1752
Cryptolestes ferrugineus, 1752
Plodia interpunctella, 1752
SADIE, 1752
Lepidoptera,
fall armyworm, 299
maize, 299
plant–herbivore interactions, 299
rootworm, 299
lesser cornstalk borer, 155, 474
burnt cane, 155
compensation, 474
CP 78-1628, 474
CP 88-1762, 474
CP 89-2143, 474
green cane, 155
sugarcane, 155
tillage, 155
lethal contraction,
Coptotermes formosanus, 1002
lethal dose, 1002
Reticulitermes flavipes, 1002
toxicity ranking, 1002
lethal dose,
Coptotermes formosanus, 1002
lethal contraction, 1002
Reticulitermes flavipes, 1002
toxicity ranking, 1002
lettuce,
mitochondrial DNA, 1280
multiplex polymerase chain reaction, 1280
Plusiinae, 1280
species identification, 1280
liberibacter, 1486, 1783
inoculation efficiency, 1486
potato, 1486, 1783
potato psyllid, 1486, 1783
psyllid density, 1783
zebra chip, 1486
zebra chip disease, 1783
life cycle, 32, 465
biological control, 32
emergence, 465
insect pest, 465
leafcutter ants, 32
native phorid species, 32
parasitism rate, 32
Sminthurus viridis, 465
soil moisture, 465
life history traits,
insecticide resistance, 2039
naled, 2039
oriental fruit fly, 2039
population dynamics, 2039
life table, 1379, 1436
Aphis gossypii, 1379
Cucumis sativus, 1379
development, 1436
plant resistance, 1379
psocid, 1436
stored products, 1436
temperature, 1436
light intensity,
infestation levels, 1675
mound size, 1675
parasitism frequencies, 1675
lipid peroxidation,
antioxidant defenses, 827
Cd^{2+}, 827
oxidative stress, 827
Propsilocerus akamusi, 827
lipids,
carbohydrates, 1229
methoxyfenozide, 1229
proteins, 1229
Spodoptera exigua, 1229
Liposcelis,
insect growth regulator, 1765
Psocoptera, 1765
pyriproxyfen, 1765
surface treatment, 1765
Liposcelis corrodens,
heat treatment, 1415
Liposcelis decolor, 1415
Rhyzopertha dominica, 1415
Tribolium castaneum, 1415
Liposcelis decolor,
heat treatment, 1415
Liposcelis corrodens, 1415
Rhyzopertha dominica, 1415
Tribolium castaneum, 1415
Liriomyza sativae,
biological invasion, 1771
insecticide susceptibility, 1771
Liriomyza trifolii, 1771
species displacement, 1771
Liriomyza trifolii,
biological invasion, 1771
insecticide susceptibility, 1771
Liriomyza sativae, 1771
species displacement, 1771
Lobesia botrana,
bunch damage, 580
Cyprus, 580
insecticides, 580
vineyard, 580
long-distance transportation,
codling moth, 814
sterile insect technique, 814
longevity, 509, 1212
Ephestia kuehniella, 1212
fecundity, 509, 1212
fumigant toxicity, 1212
phenology, 509
rearing, 509
survival, 509
lower developmental threshold, 1286, 1870
base temperature, 1870
citrus peelminer, 1286
day degrees, 1870
degree-day, 1286
rate of development, 1870
sum of effective temperatures, 1870
low pressure,
Mediterranean fruit fly, 75
melon fly, 75
oriental fruit fly, 75
phytosanitary treatment, 75
Loxosceles reclusa,
pest control, 986
pesticide, 986
scavenging, 986
lufenuron,
Adress, 1647
Ceratitis capitata, 1647
chemosterilant, 1647
SIT, 1647
Lycoriella ingenua,
control, 839
insect growth regulator, 839
mushroom sciarid fly, 839
soil drench, 839
Lygus elisus,
crop damage, 1575
Lygus lineolaris, 1575
Lygus hesperus, 1575
Physaria fendleri, 1575
Lygus lineolaris,
crop damage, 1575
Lygus elisus, 1575
Lygus hesperus, 1575
Physaria fendleri, 1575
Lygus hesperus,
crop damage, 1575
Lygus elisus, 1575
Lygus lineolaris, 1575
Physaria fendleri, 1575
Lymantria dispar,
biological invasions, 1265
IPM, 1265
pheromone release rate, 1265
pheromone-baited traps, 1265

M

Macchiademus diplopterus,
controlled atmosphere temperature treatment system, 398
heat, 398
Phlyctinus callosus, 398
postharvest, 398
Macrosteles quadrilineatus,
aster leafhopper, 1800
aster yellows, 1800
aster yellows phytoplasma, 1800
qPCR, 1800

Madeira mealybug,
chrysanthemyl 2-methylbutanoate, 823
field bioassay, 823
lavandulyl 2-methylbutanoate, 823
stereochemistry, 823
maize,
fall armyworm, 299
Lepidoptera, 299
plant–herbivore interactions, 299
rootworm, 299
malathion,
Anastrepha ludens, 452
fruit fly, 452
spinosad, 452
traps, 452
management,
Allium cepa, 1
Iris yellow spot virus, 1
Thrips tabaci, 1
mango,
biomolecules, 1991
fruit fly, 1991
graft, 1991
preharvest, 1991
manuka oil,
detection tools, 517
double-decker trap, 517
green leaf volatiles, 517
invasive forest pest, 517
mass-marking method,
elytral color polymorphism, 420
mating compatibility, 420
sterile insect technique, 420
mating compatibility,
elytral color polymorphism, 420
mass-marking method, 420
sterile insect technique, 420
mating competitiveness,
Bactrocera dorsalis, 1969
invasive species, 1969
sterile male releases, 1969
mating disruption,
competitive attraction, 939
pheromone dispenser, 939
Vitacea polistiformis, 939
mating failure,
asynchronous development, 1146
nonreproduction, 1146
resistance, 1146
Varroa destructor, 1146
mating status,
flight initiation, 2078
presence or absence of food, 2078
starvation, 2078
Tribolium castaneum, 2078
Mayetiola destructor, 792, 1384, 1393
avirulence gene, 1393
coevolution, 1393
fumigation, 792
Hessian fly, 1384
hydrogen phosphide, 792
Oulema melanopus, 792
phytohormones, 1384
plant breeding, 1393
pressure, 792
pyramiding, 1393
rice, 1384
wheat, 1384
mCry3A, 1584, 2054
Bt, 1584
Diabrotica virgifera virgifera, 1584, 2054
eCry3.1Ab, 1584
insect resistance management, 2054
phenology, 2054
resistance management, 1584
mealworms,
bioassay, 2095
Ixodes scapularis, 2095
Metarhizium brunneum, 2095
Tenebrio molitor, 2095
mealybugs,
detection, 555
ornamental plants, 555
pheromone trap, 555
sampling, 555
mechanism,
Cydia molesta, 636
insecticide resistance, 636
monitoring, 636
target mutation, 636
Mediterranean fruit fly, 75, 188, 1349, 1856
bait station, 1856
behavior, 1856
chemical control, 1349
disinfection, 188
eggs, 188
esterases, 1349
integrated pest management, 1856
low pressure, 75
melon fly, 75
oriental fruit fly, 75
phytosanitary treatment, 75
quaternary ammonium, 188
rainfastness, 1856
rearing, 188
resistance management, 1349
synergist, 1349
Megachile rotundata,
alternative pollinator, 1162
cold storage, 1162
fluctuating thermal regime, 1162
fluctuation thermal regime, 1162
melon fly,
low pressure, 75
Mediterranean fruit fly, 75
oriental fruit fly, 75
phytosanitary treatment, 75
menthol,
1-octen-3-ol, 1633
borneol, 1633
repellency, 1633
volatiles, 1633
metabolic resistance,
Alphitobius diaperinus, 629
cross-resistance, 629
darkling beetle, 629
piperonyl butoxide, 629
Metarhizium brunneum,
bioassay, 2095
Ixodes scapularis, 2095
mealworms, 2095
Tenebrio molitor, 2095
methionine,
biorational, 1986
larvicide, 1986
methoxyfenozide,
carbohydrates, 1229
lipids, 1229
proteins, 1229
Spodoptera exigua, 1229
methyl anthranilate,
acetic acid, 1195
grape juice, 1195
Mexican fruit fly, 1195
propylene glycol, 1195
methyl bromide,
Agrilus, 69
Fraxinus, 69
fumigation, 69
quarantine, 69
methyl bromide alternative,
heat tolerance, 317
heat treatment, 317
methyl salicylate,
herbivore-induced plant volatile, 115
lacewing, 115
soybean aphid, 115
syrphid fly, 115
Mexican fruit fly,
acetic acid, 1195
grape juice, 1195
methyl anthranilate, 1195
propylene glycol, 1195
microsatellites,
Bactrocera cucurbitae, 1065
gene flow, 1065
genetic structure, 1065
quarantine pest, 1065
MIR604,
Bacillus thuringiensis, 1045
Diabrotica virgifera virgifera, 1045
toxicity assay, 1045
mirid,
biological control, 1742
integrated pest management, 1742
monitoring, 1742
pear psyllid, 1742
mite resistance,
Apis mellifera, 745
breeding, 745
Varroa destructor, 745
mitochondrial DNA,
lettuce, 1280
multiplex polymerase chain reaction, 1280

Plusiinae, 1280
species identification, 1280
mitochondrial population genetics,
Epiphyas postvittana, 920
haplotype, 920
invasive species, 920
origin, 920
modeling,
aeration, 702
insects, 702
rice, 702
storage, 702
modified atmosphere,
carbon dioxide, 1774
environment, 1774
mortality, 1774
Varroa destructor, 1774
MON863,
Bt resistance, 1038
Diabrotica virgifera virgifera, 1038
insect resistance management, 1038
monitoring, 173, 211, 274, 636, 1093, 1258, 1613, 1742
aggregation, 1613
almonds, 211
Amyelois transitella, 211
ash canopy, 173
attractant, 1258
bed bug, 274
biological control, 1742
Cerambycidae, 1258
Cimex lectularius, 274
Cydia molesta, 636
dairy, 1093
dry ice, 274
green prism traps, 173
house fly, 1093
insecticide resistance, 636
integrated pest management, 1613, 1742
IPM, 1093
mechanism, 636
mirid, 1742
oviposition traps, 211
pear psyllid, 1742
phoebe oil, 173
pistachios, 211
reduced-risk, 1613
semiochemicals, 1613
surveillance, 1093
target mutation, 636
trap design, 1258
traps, 274
woodborer, 1258
Z-3-hexenol, 173
monitoring station,
feeding rates, 1024
seasonal trends, 1024
subterranean termite, 1024
monitors,
Heterotermes aureus, 1339
subterranean termites, 1339
trapping, 1339
Monochamus alternatus,
2-(undecyloxy)-ethanol, 1592
aggregation pheromone, 1592
ethanol, 1592
α-pinene, 1592
Monochamus titillator, 1245, 1934
Buprestidae, 1934
Cerambycidae, 1934
Curculionidae, 1934
ethanol, 1245
ipsdienol, 1245
ipsenol, 1245
α-pinene, 1245
Scolytinae, 1934
Monomorium chinense,
interspecific competition, 614
Pheidole fervens, 614
Solenopsis invicta, 614
Solenopsis invicta virus-1, 614
monoterpenes,
Blattella germanica, 595
essential oils, 595
Eucalyptus, 595
knockdown, 595
mortality,
carbon dioxide, 1774
environment, 1774
modified atmosphere, 1774
Varroa destructor, 1774
mosquito, 1369, 1816
Aedes aegypti, 1369
amide, 1369
B. thuringiensis subsp. *Israelensis*, 1816
Bacillus sphaericus, 1816
behavioral response, 1369
biopesticides, 1816
Coffee husk waste, 1816
repellent, 1369
mosquito control,
larval development, 266
pupation rate, 266
sterile insect technique, 266
survival rate, 266
mound size,
infestation levels, 1675
light intensity, 1675
parasitism frequencies, 1675
mRNA expression,
bacterial challenge, 2046
glutathione S-transferase, 2046
homology modeling, 2046
insecticides, 2046
multiplex polymerase chain reaction,
lettuce, 1280
mitochondrial DNA, 1280
Plusiinae, 1280
species identification, 1280
Musca domestica,
Muscidifurax, 1108
population dynamics, 1108
Spalangia, 1108
Stomoxys calcitrans, 1108
Muscidifurax,
Musca domestica, 1108
population dynamics, 1108
Spalangia, 1108
Stomoxys calcitrans, 1108
mushroom sciarid fly,
control, 839
insect growth regulator, 839
Lycoriella ingenua, 839
soil drench, 839

N

naled,
insecticide resistance, 2039
life history traits, 2039
oriental fruit fly, 2039
population dynamics, 2039
Nantucket pine tip moth,
fipronil, 1272
imidacloprid, 1272
pales weevil, 1272
systemic insecticides, 1272
native elm bark beetle,
American elm, 889
bioassay, 889
Dutch elm disease, 889
efficacy, 889
native phorid species,
biological control, 32
leafcutter ants, 32
life cycle, 32
parasitism rate, 32
natural enemy, 845, 1125
integrated pest management, 1125
IPM, 845
pheromone mating disruption, 1125
plant volatiles, 1125
population dynamics, 845
natural lure,
Drosophila, 1018
trap, 1018
navel orangeworm, 532, 965
almonds, 965
Amyelois transitella, 965
degree-days, 532
development, 532
hull split, 965
pistachios, 532
shell seal, 965
survival, 532
nectar production,
diurnal pollination, 149
female reproductive success, 149
Jatropha curcas, 149
nocturnal pollination, 149
neonicotinoid, 205, 405, 1942
Anoplophora glabripennis, 1942
bioassays residual activity, 405
Blissus occiduus, 205
buffalograss, 205
chinch bug, 205
chronic exposure, 405
fecundity, 1942

invasive species, 1942
LC_{50}, 205
Otiorhynchus sulcatus, 405
sublethal effect, 1942
neonicotinoid insecticides,
reproductively mature flies, 194
Rhagoletis indifferens, 194
spinosad bait, 194
sucrose, 194
Neotephritis finalis,
planting date, 1236
Suleima helianthana, 1236
sunflower, 1236
sunflower bud moth, 1236
Neotyphodium,
alkaloid, 665
corn flea beetle, 665
endophyte, 665
insect resistance, 665
Nephotettix cincticeps,
Laodelphax striatellus, 1877
Nilaparvata lugens, 1877
pymetrozine, 1877
Sogatella furcifera, 1877
Nicaragua,
Brassicae oleracea, 782
Diadegma insulare, 782
Plutella xylostella, 782
Polybia sp., 782
Nilaparvata lugens, 243, 1364, 1877
biochemical mechanism, 243
bisultap resistant strain, 243
butene-fipronil, 1364
cross-resistance, 243, 1364
ethiprole, 1364
fipronil, 1364
Laodelphax striatellus, 1877
Nephotettix cincticeps, 1877
pymetrozine, 1877
Sogatella furcifera, 1877
nitrogen fertilizer,
economic threshold, 1550
field pea, 1550
pea leaf weevil, 1550
yield loss, 1550
Noctuidae,
Bacillus thuringiensis kurstaki and *aizawai*, 41
black cutworm, 41
nocturnal pollination,
diurnal pollination, 149
female reproductive success, 149
Jatropha curcas, 149
nectar production, 149
nonchemical control, 325, 1423
Acari, 1423
Acaridae, 1423
disinfestation, 1423
integrated pest management, 325
stored grain, 325
supercooling point, 1423
nonrepellent,
Coptotermes formosanus, 2025
delayed action, 2025
substrate type, 2025
termiticide, 2025
nonreproduction,
asynchronous development, 1146
mating failure, 1146
resistance, 1146
Varroa destructor, 1146
nontarget impacts,
Aphis glycines, 1824
Entomophthorales, 1824
Glycine max, 1824
Rag1, 1824
nontargets,
Folsomia candida, 1892
risk assessment, 1892
transgenic Bt rice, 1892
nootkatone,
Amblyomma americanum, 659
carvacrol, 659
EcoSMART, 659
Ixodes scapularis, 659
North America,
Contarinia nasturtii, 709
integrated pest management, 709
invasive species, 709
nymphoid,
Cryptotermes, 622
de-alates, 622
sex determination, 622
sex ratio, 622

O

observational studies,
causal inference, 331
ecoinformatics, 331
economic injury level, 331
statistical power, 331
1-octen-3-ol,
borneol, 1633
menthol, 1633
repellency, 1633
volatiles, 1633
Odontotermes formosanus,
Klason lignin, 862
termite feeding, 862
wood extractive, 862
wood hardness, 862
odor recognition,
bait trap station, 225
detection dog, 225
fire ant detection, 225
red imported fire ant, 225
oil,
alcohol, 1680
cockroach, 1680
synergism, 1680
toxicity, 1680
Oligonycus aceris,
economic analysis, 1960
insecticide coverage, 1960
secondary pest outbreak, 1960
Xylosandrus crassiusculus, 1960
Ooencyrtus nezarae,
cold storage, 1833
conservational biological control, 1833
Gryon japonicum, 1833
stink bug, 1833
oothecal parasitoid,
biological control, 2031
cockroach control, 2031
host age preference, 2031
Periplaneta americana, 2031
Ophelimus maskelli,
damage, 180
gall density, 180
kaolin, 180
organic,
botanical insecticides, 1979
damage, 1979
grapefruit, 1979
Kelly's citrus thrips, 1979
Pezothrips kellyanus, 1979
organic cultivation,
coffee, 1909
coffee insect pests, 1909
conventional cultivation, 1909
semiochemical, 1909
organophosphate,
adjuvant, 1309
insecticide management, 1309
kairomone, 1309
pyrethroid, 1309
oriental fruit fly, 75, 2039
insecticide resistance, 2039
life history traits, 2039
low pressure, 75
Mediterranean fruit fly, 75
melon fly, 75
naled, 2039
phytosanitary treatment, 75
population dynamics, 2039
oriental fruit moth,
baseline susceptibility, 1950
codling moth, 1950
reduced-risk insecticides, 1950
origin,
Epiphyas postvittana, 920
haplotype, 920
invasive species, 920
mitochondrial population genetics, 920
ornamental plants,
detection, 555
mealybugs, 555
pheromone trap, 555
sampling, 555
Osmia lignaria,
bee dispersal, 750
bee retention, 750
solitary bees, 750
Ostrinia nubilalis,
Bacillus thuringiensis, 1476
Helicoverpa zea, 1476
resistance management, 1476

Otiorhynchus sulcatus,
bioassays residual activity, 405
chronic exposure, 405
neonicotinoids, 405
Oulema melanopus,
fumigation, 792
hydrogen phosphide, 792
Mayetiola destructor, 792
pressure, 792
out-hive traps,
Aethina tumida, 26
in-hive traps, 26
polyvinyl chloride trap, 26
white trap, 26
ovicidal activity,
Epiphyas postvittana, 220
petroleum oils, 220
reduced risk pesticides, 220
oviposition, 753, 1622
Aethina tumida, 753
Allium cepa, 1622
cultural control, 1622
Delia antiqua, 1622
diet, 753
planting date, 1622
survivorship, 753
temperature, 753
oviposition traps,
almonds, 211
Amyelois transitella, 211
monitoring, 211
pistachios, 211
oxidative stress, 101, 827
antioxidant defenses, 827
biomarker, 101
Cd^{2+}, 827
glutathione transferase, 101
lipid peroxidation, 827
phoxim, 101
Propsilocerus akamusi, 827
silkworm, 101
oxygen,
phosphine fumigation, 1455
postharvest pest control, 1455
stored product, 1455
ozone, 308, 353
chalkbrood, 353
foulbrood, 353
honey bees, 353
insect control, 308
pathogens, 353
psocids, 308
stored grain beetles, 308
stored wheat, 308

P

P. pruinosa,
common eastern bumble bee, 1153
European honey bee, 1153
foraging behavior, 1153
pumpkin, 1153
pales weevil,
fipronil, 1272
imidacloprid, 1272
Nantucket pine tip moth, 1272
systemic insecticides, 1272
paper consumption,
Coptotermes formosanus, 232
Reticulitermes speratus, 232
termite lipid content, 232
termite survival, 232
parasitism,
Dasineura mali, 1640
phenology, 1640
Platygaster demades, 1640
synchronization, 1640
parasitism frequencies,
infestation levels, 1675
light intensity, 1675
mound size, 1675
parasitism rate,
biological control, 32
leafcutter ants, 32
life cycle, 32
native phorid species, 32
parasitoid rearing,
Agrilus planipennis, 379
biocontrol, 379
Spathius agrili, 379
path analysis,
above and belowground interactions, 1568
aphid colonization, 1568
Aphid glycines, 1568
Heterodera glycines, 1568
pathogens,
chalkbrood, 353
foulbrood, 353
honey bees, 353
ozone, 353
pea leaf weevil,
economic threshold, 1550
field pea, 1550
nitrogen fertilizer, 1550
yield loss, 1550
peach,
Heterorhabditis, 47
Prunus persicae, 47
Steinernema carpocapsae, 47
Synanthedon pictipes, 47
pear psyllid,
biological control, 1742
integrated pest management, 1742
mirid, 1742
monitoring, 1742
Pentatomidae,
chemical control, 137
egg efficacy, 137
Scelionidae, 137
Periplaneta americana,
biological control, 2031
cockroach control, 2031
host age preference, 2031
oothecal parasitoid, 2031
persistence,
chlorantraniliprole, 990
indoxacarb, 990
R. flavipes, 990
termiticide, 990
pest, 764, 1561
bioenergy, 1561
defoliator, 1561
ethanol, 1561
invasive insect species, 764
Kalman filter, 764
resistance, 1561
state-space model, 764
pest control, 601, 986
Arachnida, 601
Holocnemus pluchei, 601
Loxosceles reclusa, 986
pesticide, 986
scavenging, 986
urban entomology, 601
pest management,
3,5-dimethyldodecanoic acid, 933
pheromone, 933
pest risk analysis,
host range, 1514
host specificity, 1514
invasive species, 1514
Tortricidae, 1514
pesticide,
Loxosceles reclusa, 986
pest control, 986
scavenging, 986
petroleum oils,
Epiphyas postvittana, 220
ovicidal activity, 220
reduced risk pesticides, 220
Pezothrips kellyanus,
botanical insecticides, 1979
damage, 1979
grapefruit, 1979
Kelly's citrus thrips, 1979
organic, 1979
Pheidole fervens,
interspecific competition, 614
Monomorium chinense, 614
Solenopsis invicta, 614
Solenopsis invicta virus-1, 614
phenology, 509, 1640, 2054
Dasineura mali, 1640
Diabrotica virgifera virgifera, 2054
fecundity, 509
insect resistance management, 2054
longevity, 509
mCry3A, 2054
parasitism, 1640
Platygaster demades, 1640
rearing, 509
survival, 509
synchronization, 1640
pheromone,
3,5-dimethyldodecanoic acid, 933
pest management, 933
pheromone dispenser,
competitive attraction, 939
mating disruption, 939
Vitacea polistiformis, 939

pheromone mating disruption,
integrated pest management, 1125
natural enemies, 1125
plant volatiles, 1125
pheromone release rate,
biological invasions, 1265
IPM, 1265
Lymantria dispar, 1265
pheromone-baited traps, 1265
pheromone trap,
detection, 555
mealybugs, 555
ornamental plants, 555
sampling, 555
pheromone trap survey,
forest defoliator, 494
invasive species, 494
pheromone-baited traps,
biological invasions, 1265
IPM, 1265
Lymantria dispar, 1265
pheromone release rate, 1265
Phlyctinus callosus,
controlled atmosphere temperature treatment system, 398
heat, 398
Macchiademus diplopterus, 398
postharvest, 398
phoebe oil,
ash canopy, 173
green prism traps, 173
monitoring, 173
Z-3-hexenol, 173
phosphine fumigation,
oxygen, 1455
postharvest pest control, 1455
stored product, 1455
photosynthesis,
Empoasca fabae, 1316
host plant expansion, 1316
transpiration, 1316
phoxim,
biomarker, 101
glutathione transferase, 101
oxidative stress, 101
silkworm, 101
Phyllotreta spp.,
canola, 125
flea beetle feeding, 125
host plant resistance, 125
trichomes, 125
Physaria fendleri,
crop damage, 1575
Lygus elisus, 1575
Lygus lineolaris, 1575
Lygus hesperus, 1575
phytohormones,
Hessian fly, 1384
Mayetiola destructor, 1384
rice, 1384
wheat, 1384
phytoplasma transmission,
broad bean, 360
chrysanthemum yellows, 360
Flavescence dorée, 360
leafhopper, 360
phytosanitary, 63, 1180
Anastrepha ludens, 63
Bactrocera invadens, 1180
cold disinfestation, 1180
commodity treatment, 63, 1180
quarantine, 63, 1180
phytosanitary regulation,
Ceratitis capitata, 1174
cold disinfestation, 1174
commodity treatment, 1174
quarantine, 1174
phytosanitary treatment, 21, 75, 1509, 1851
American serpentine leafminer, 1851
Bactrocera cucurbitae, 1509
Bactrocera latifrons, 1509
cold treatment, 21
heat treatment, 21
hot water immersion, 21
ionizing radiation, 1851
low pressure, 75
Mediterranean fruit fly, 75
melon fly, 75
oriental fruit fly, 75
quarantine pest, 1509
quarantine treatment, 21, 1851
X-ray irradiation, 1509
α-pinene, 574, 1245, 1592
aggregation pheromone, 1592
ambrosia beetles, 574
ethanol, 574, 1245, 1592
ipsdienol, 1245
ipsenol, 1245
Monochamus alternatus, 1592
Monochamus titillator, 1245
Scolytinae, 574
2-(undecyloxy)-ethanol, 1592
piperonyl butoxide,
Alphitobius diaperinus, 629
cross-resistance, 629
darkling beetle, 629
metabolic resistance, 629
pistachios, 211, 532
almonds, 211
Amyelois transitella, 211
degree-days, 532
development, 532
monitoring, 211
navel orangeworm, 532
oviposition traps, 211
survival, 532
pitfall trapping,
insect vision, 1430
LED, 1430
stored product, 1430
Planococcus ficus,
Ampelovirus, 1480
Closteroviridae, 1480
semipersistent, 1480
plant breeding,
avirulence gene, 1393
coevolution, 1393
Mayetiola destructor, 1393
pyramiding, 1393
plant bug,
cotton, 685
sampling, 685
sweep net, 685
plant resistance, 1379, 2073
Aphis gossypii, 1379
Blissus occiduus, 2073
Buchloe dactyloides, 2073
Cucumis sativus, 1379
life table, 1379
turfgrass, 2073
plant volatiles,
integrated pest management, 1125
natural enemies, 1125
pheromone mating disruption, 1125
plant–herbivore interactions,
fall armyworm, 299
Lepidoptera, 299
maize, 299
plant–herbivore interactions, 299
rootworm, 299
planting date, 1236, 1622
Allium cepa, 1622
cultural control, 1622
Delia antiqua, 1622
Neotephritis finalis, 1236
oviposition, 1622
Suleima helianthana, 1236
sunflower, 1236
sunflower bud moth, 1236
Platygaster demades,
Dasineura mali, 1640
parasitism, 1640
phenology, 1640
synchronization, 1640
Plodia interpunctella, 693, 1752
biological control, 693
Blattisocius tarsalis, 1752
Cryptolestes ferrugineus, 1752
intraspecific competition, 693
Lepidoglyphus destructor, 1752
SADIE, 1752
Sitophilus oryzae, 693
Plusiinae,
lettuce, 1280
mitochondrial DNA, 1280
multiplex polymerase chain reaction, 1280
species identification, 1280
Plutella xylostella,
Brassicae oleracea, 782
Diadegma insulare, 782
Nicaragua, 782
Polybia sp., 782
poinsettia,
Bemisia tabaci, 548
Euphorbia pulcherrima, 548
sweetpotato whitefly, 548
Polistinae,
diversity, 1120
Eucalyptus, 1120

pollination limitation,
agriculture, 107
Bombus terrestris, 107
Rubus idaeus, 107
Scotland, 107
Polybia sp.,
Brassicae oleracea, 782
Diadegma insulare, 782
Nicaragua, 782
Plutella xylostella, 782
polyvinyl chloride trap,
Aethina tumida, 26
in-hive traps, 26
out-hive traps, 26
white trap, 26
Popillia japonica,
predation, 2009
secondary pest outbreak, 2009
Solenopsis, 2009
poplar petiolegall aphid,
apterae, 1116
extraction, 1116
sampling, 1116
spatial distribution, 1116
population dynamics, 845, 1108, 2039
insecticide resistance, 2039
IPM, 845
life history traits, 2039
Musca domestica, 1108
Muscidifurax, 1108
naled, 2039
natural enemy, 845
oriental fruit fly, 2039
Spalangia, 1108
Stomoxys calcitrans, 1108
population growth, 799, 2087
biological control, 799
developmental period, 2087
fecundity, 799
flour, 2087
intrinsic rate of increase, 799
starch, 2087
temperature threshold, 799
Tribolium castaneum, 2087
population growth rate,
development rate, 1864
fecundity, 1864
thermal requirements, 1864
postharvest,
controlled atmosphere temperature treatment system, 398
heat, 398
Macchiademus diplopterus, 398
Phlyctinus callosus, 398
postharvest pest control,
oxygen, 1455
phosphine fumigation, 1455
stored product, 1455
potato, 586, 1075, 1486, 1783
accession, 1075
aphid, 1075
breeding, 1075
inoculation efficiency, 1486
insecticides, 586
IPM, 586
liberibacter, 1486, 1783
potato psyllid, 586, 1486, 1783
psyllid density, 1783
resistance, 1075
zebra chip, 586, 1486
zebra chip disease, 1783
potato psyllid, 586, 1138, 1486, 1783
inoculation efficiency, 1486
insecticides, 586
insecticide spraying, 1138
integrated pest management, 1138
IPM, 586
liberibacter, 1486, 1783
potato, 586, 1486, 1783
psyllid density, 1783
row crops, 1138
water-sensitive spray cards, 1138
zebra chip, 586, 1486
zebra chip disease, 1783
precision,
accuracy, 1323
efficiency, 1323
Frankliniella occidentalis, 1323
sampling, 1323
predation,
Popillia japonica, 2009
secondary pest outbreak, 2009
Solenopsis, 2009
predators, 773, 868
beneficial insects, 773
eastern hemlock, 868
egg parasitoids, 773
fertilizer, 868
hemlock woolly adelgid, 868
imidacloprid, 868
laboratory bioassay, 773
systemic insecticides, 773
predatory mites,
Calepitrimerus vitis, 970
integrated pest management, 970
sublethal effects, 970
toxicity, 970
preharvest,
biomolecules, 1991
fruit fly, 1991
graft, 1991
mango, 1991
presence or absence of food,
flight initiation, 2078
mating status, 2078
starvation, 2078
Tribolium castaneum, 2078
pressure,
fumigation, 792
hydrogen phosphide, 792
Mayetiola destructor, 792
Oulema melanopus, 792
prism trap,
emerald ash borer, 901
funnel trap, 901
trap color, 901
visual cues, 901
Propsilocerus akamusi,
antioxidant defenses, 827
Cd^{2+}, 827
lipid peroxidation, 827
oxidative stress, 827
propylene glycol,
acetic acid, 1195
grape juice, 1195
methyl anthranilate, 1195
Mexican fruit fly, 1195
proteins,
carbohydrates, 1229
lipids, 1229
methoxyfenozide, 1229
Spodoptera exigua, 1229
Prunus persicae,
Heterorhabditis, 47
peach, 47
Steinernema carpocapsae, 47
Synanthedon pictipes, 47
Psidium guajava,
Anastrepha, 1204
competitive displacement, 1204
host status, 1204
infestation patterns, 1204
psocid, 308, 1436
development, 1436
insect control, 308
life table, 1436
ozone, 308
stored grain beetles, 308
stored products, 1436
stored wheat, 308
temperature, 1436
Psocoptera, 1720, 1765
DNA barcoding, 1720
insect growth regulator, 1765
internal transcribed spacer, 1720
Liposcelis, 1765
pyriproxyfen, 1765
species identification, 1720
surface treatment, 1765
psyllid density,
liberibacter, 1783
potato, 1783
potato psyllid, 1783
zebra chip disease, 1783
Psyttalia concolor,
fipronil, 1541
imidacloprid, 1541
side effects, 1541
pumpkin,
common eastern bumble bee, 1153
European honey bee, 1153
foraging behavior, 1153
P. pruinosa, 1153
pupation rate,
larval development, 266
mosquito control, 266
sterile insect technique, 266
survival rate, 266
pymetrozine,
Laodelphax striatellus, 1877
Nephotettix cincticeps, 1877

Nilaparvata lugens, 1877
Sogatella furcifera, 1877
pyramiding,
avirulence gene, 1393
coevolution, 1393
Mayetiola destructor, 1393
plant breeding, 1393
pyrethroid, 250, 1309
adjuvant, 1309
insecticide management, 1309
kairomone, 1309
organophosphate, 1309
Stomoxys calcitrans, 250
voltage-gated sodium channel, 250
pyriproxyfen,
insect growth regulator, 1765
Liposcelis, 1765
Psocoptera, 1765
surface treatment, 1765

Q

qPCR,
aster leafhopper, 1800
aster yellows, 1800
aster yellows phytoplasma, 1800
Macrosteles quadrilineatus, 1800
quarantine, 63, 69, 414, 1174, 1180
Agrilus, 69
Anastrepha ludens, 63
Bactrocera invadens, 1180
Ceratitis capitata, 1174
citral, 414
cold disinfestation, 1174, 1180
commodity treatment, 63, 1174, 1180
common thyme, 414
essential oils, 414
Fraxinus, 69
fumigation, 69
lemongrass, 414
methyl bromide, 69
phytosanitary, 63, 1180
phytosanitary regulation, 1174
quarantine pest, 1065, 1509
Bactrocera cucurbitae, 1065, 1509
Bactrocera latifrons, 1509
gene flow, 1065
genetic structure, 1065
microsatellites, 1065
phytosanitary treatment, 1509
X-ray irradiation, 1509
quarantine treatment, 21, 717, 1851
American serpentine leafminer, 1851
cold treatment, 21
heat treatment, 21
hot water immersion, 21
International Plant Protection Convention, 717
International Standard for Phytosanitary Measures No. 15, 717
ionizing radiation, 1851
phytosanitary treatment, 21, 1851
quaternary ammonium,
disinfection, 188
eggs, 188
Mediterranean fruit fly, 188
rearing, 188

R

R. flavipes,
chlorantraniliprole, 990
indoxacarb, 990
persistence, 990
termiticide, 990
radiation,
Epiphyas postvittana, 1301
eradication, 1301
sterile insect technique, 1301
Tortricidae, 1301
Rag1,
Aphis glycines, 1824
Entomophthorales, 1824
Glycine max, 1824
nontarget impacts, 1824
rainfastness, 1656, 1856
bait station, 1856
behavior, 1856
grapes, 1656
integrated pest management, 1856
Japanese beetle, 1656
Mediterranean fruit fly, 1856
residual activity, 1656
residue profile, 1656
rate of development,
base temperature, 1870
day degrees, 1870
lower developmental threshold, 1870
sum of effective temperatures, 1870
real-time polymerase chain reaction,
barcodes, 1706
internal transcribed spacer, 1706
Tortricidae, 1706
rearing, 188, 509
disinfection, 188
eggs, 188
fecundity, 509
longevity, 509
Mediterranean fruit fly, 188
phenology, 509
quaternary ammonium, 188
survival, 509
rearing density,
flight initiation, 443
food deprivation, 443
red flour beetle, 443
Tribolium castaneum, 443
red flour beetle,
flight initiation, 443
food deprivation, 443
rearing density, 443
Tribolium castaneum, 443
red imported fire ant,
bait trap station, 225
detection dog, 225
fire ant detection, 225
odor recognition, 225
reduced risk pesticides,
Epiphyas postvittana, 220
ovicidal activity, 220
petroleum oils, 220
reduced-risk,
aggregation, 1613
integrated pest management, 1613
monitoring, 1613
semiochemicals, 1613
reduced-risk insecticides,
baseline susceptibility, 1950
codling moth, 1950
oriental fruit moth, 1950
refuge,
biological control, 343
Bt corn, 343
insect resistance management, 343
secondary pests, 343
relative humidity,
C. modoc, 740
C. vicinus, 740
carpenter ants, 740
colony founding, 740
repellency,
1-octen-3-ol, 1633
borneol, 1633
menthol, 1633
volatiles, 1633
repellent,
Aedes aegypti, 1369
amide, 1369
behavioral response, 1369
mosquito, 1369
repellent testing,
Amblyomma americanum, 2062
Ixodes scapularis, 2062
reproductively mature flies,
neonicotinoid insecticides, 194
Rhagoletis indifferens, 194
spinosad bait, 194
sucrose, 194
resampling software,
benefit-cost ratio, 1900
binomial sampling, 1900
western bean cutworm, 1900
residual activity,
grapes, 1656
Japanese beetle, 1656
rainfastness, 1656
residue profile, 1656
residue profile,
grapes, 1656
Japanese beetle, 1656
rainfastness, 1656
residual activity, 1656
resistance, 566, 673, 1075, 1146, 1406, 1561, 1730, 2068
accession, 1075
aphid, 1075
asynchronous development, 1146
Bacillus thuringiensis, 673

Bactrocera cucurbitae, 566
Bactrocera dorsalis, 566
bioenergy, 1561
breeding, 1075
Ceratitis capitata, 566
Cnaphalocrocis medinalis, 1730
cotton, 673
cowpea trypsin inhibitor, 673
Cry1C*, 1730
Cry2A*, 1730
defoliator, 1561
EPG, 2068
ethanol, 1561
feeding behavior, 2068
Helicoverpa armigera, 673
interaction, 1406
mating failure, 1146
nonreproduction, 1146
pest, 1561
potato, 1075
rice, 1730
soybean aphids, 2068
Vaccinium spp., 566
Varroa destructor, 1146
wheat, 1406
wheat curl mite, 1406
wheat streak mosaic virus, 1406
resistance management, 806, 1343, 1349, 1476, 1584
Bacillus thuringiensis, 1476
blended seed refuge, 806
Bt, 1584
chemical control, 1349
Diabrotica virgifera virgifera, 1584
eCry3.1Ab, 1584
esterases, 1349
European corn borer, 806
feeding behavior, 806
generating susceptible strain, 1343
Helicoverpa zea, 1476
insecticide resistance, 1343
mCry3A, 1584
Mediterranean fruit fly, 1349
Ostrinia nubilalis, 1476
Spodoptera exigua, 1343
synergist, 1349
transgenic corn, 806
resistance monitoring,
bioassays, 1031
Grapholita molesta, 1031
Reticulitermes flavipes, 425, 1002
cations, 425
Coptotermes formosanus, 1002
foraging, 425
lethal contraction, 1002
lethal dose, 1002
soil, 425
termite, 425
toxicity ranking, 1002
Reticulitermes speratus,
Coptotermes formosanus, 232
paper consumption, 232
termite lipid content, 232
termite survival, 232
Rhagoletis indifferens,
neonicotinoid insecticides, 194
reproductively mature flies, 194
spinosad bait, 194
sucrose, 194
Rhopalosiphum padi,
aphids, 1167
barley yellow dwarf, 1167
cereal yellow dwarf, 1167
virus vectors, 1167
Rhyzopertha dominica,
heat treatment, 1415
Liposcelis corrodens, 1415
Liposcelis decolor, 1415
Tribolium castaneum, 1415
rice, 702, 1384, 1730
aeration, 702
Cnaphalocrocis medinalis, 1730
Cry1C*, 1730
Cry2A*, 1730
Hessian fly, 1384
insects, 702
Mayetiola destructor, 1384
modeling, 702
phytohormones, 1384
resistance, 1730
storage, 702
wheat, 1384
risk assessment,
Folsomia candida, 1892
nontargets, 1892
transgenic Bt rice, 1892
rootworm,
fall armyworm, 299
Lepidoptera, 299
maize, 299
plant–herbivore interactions, 299
row crops,
insecticide spraying, 1138
integrated pest management, 1138
potato psyllids, 1138
water-sensitive spray cards, 1138
Rubus idaeus,
agriculture. 107
Bombus terrestris, 107
pollination limitation, 107
Scotland, 107
Russian wheat aphid,
biotype, 1736
Diuraphis noxia, 1736
host plant resistance, 1736
Rutaceae,
albedo, 54
flavedo, 54
host status, 54
Tephritidae, 54

S

28S,
cytochrome C oxidase I, 279
Graphocephala cythura, 279
Graphocephala flavovittata, 279
sharpshooters, 279
SADIE,
Blattisocius tarsalis, 1752
Cryptolestes ferrugineus, 1752
Lepidoglyphus destructor, 1752
Plodia interpunctella, 1752
sampling, 555, 685, 947, 1116, 1323
accuracy, 1323
ants, 947
aphid, 947
apterae, 1116
Banana bunchy top virus, 947
cotton, 685
detection, 555
efficiency, 1323
extraction, 1116
Frankliniella occidentalis, 1323
mealybugs, 555
ornamental plants, 555
pheromone trap, 555
plant bug, 685
poplar petiolegall aphid, 1116
precision, 1323
spatial distribution, 1116
sweep net, 685
within-plant distribution, 947
sap beetles,
common smut, 956
corn earworm, 956
edible oils, 956
husk coverage, 956
sassafras,
exotic, 501
invasive, 501
Lauraceae, 501
laurel wilt, 501
scavenging,
Loxosceles reclusa, 986
pest control, 986
pesticide, 986
Scelionidae,
chemical control, 137
egg efficacy, 137
Pentatomidae, 137
Scolytinae, 574, 1934
ambrosia beetles, 574
Buprestidae, 1934
Cerambycidae, 1934
Curculionidae, 1934
ethanol, 574
Monochamus titillator, 1934
α-pinene, 574
Scotland,
agriculture, 107
Bombus terrestris, 107
pollination limitation, 107
Rubus idaeus, 107
seasonal trends,
feeding rates, 1024
monitoring station, 1024
subterranean termite, 1024
seasonal variation,
enantiomers, 895
Ips avulsus, 895
ipsdienol, 895
semiochemical context, 895

secondary pest, 343, 607
baiting, 607
biological control, 343
bistrifluron, 607
Bt corn, 343
colony elimination, 607
Globitermes sulphureus, 607
insect resistance management, 343
refuge, 343
secondary pest outbreak, 1496, 1960, 2009
biological control, 1496
economic analysis, 1960
insecticide coverage, 1960
integrated pest management, 1496
Oligonycus aceris, 1960
Popillia japonica, 2009
predation, 2009
Solenopsis, 2009
Tetranychidae, 1496
tree nuts, 1496
Xylosandrus crassiusculus, 1960
selecron,
Aleurothrixus floccosus, 1329
citrus, 1329
cyhalothrin, 1329
woolly whiteflies, 1329
semiochemical, 1613, 1909
aggregation, 1613
coffee, 1909
coffee insect pests, 1909
conventional cultivation, 1909
integrated pest management, 1613
monitoring, 1613
organic cultivation, 1909
reduced-risk, 1613
semiochemical context,
enantiomers, 895
Ips avulsus, 895
ipsdienol, 895
seasonal variation, 895
semipersistent,
Ampelovirus, 1480
Closteroviridae, 1480
Planococcus ficus, 1480
Sesamia nonagrioides,
Bt corn, 853
development, 853
ecdysteroids, 853
juvenile hormone, 853
sex determination,
Cryptotermes, 622
de-alates, 622
nymphoid, 622
sex ratio, 622
sex pheromone,
citrus leafminer, 540
SPLAT, 540
(Z,Z)-7,11-hexadecadienal, 540
(Z,Z,*E*)-7,11,13-hexadecatrienal, 540
sex ratio,
Cryptotermes, 622
de-alates, 622
nymphoid, 622
sex determination, 622
sharpshooter, 279, 367
almond leaf scorch, 367
almonds, 367
cytochrome C oxidase I, 279
Draeculacephala minerva, 367
Graphocephala cythura, 279
Graphocephala flavovittata, 279
28S, 279
vector management, 367
shell seal,
almonds, 965
Amyelois transitella, 965
hull split, 965
navel orangeworm, 965
side effects,
fipronil, 1541
imidacloprid, 1541
Psyttalia concolor, 1541
silkworm,
biomarker, 101
glutathione transferase, 101
oxidative stress, 101
phoxim, 101
SIT,
Adress, 1647
Ceratitis capitata, 1647
chemosterilant, 1647
lufenuron, 1647
Sitobion avenae,
aphid biotype, 1080
wheat, 1080
wheat resistance to aphid, 1080
Sitophilus oryzae,
biological control, 693
intraspecific competition, 693
Plodia interpunctella, 693
Sminthurus viridis,
emergence, 465
insect pest, 465
life cycle, 465
soil moisture, 465
Sogatella furcifera,
Laodelphax striatellus, 1877
Nephotettix cincticeps, 1877
Nilaparvata lugens, 1877
pymetrozine, 1877
soil,
cations, 425
foraging, 425
Reticulitermes flavipes, 425
termite, 425
soil drench,
control, 839
insect growth regulator, 839
Lycoriella ingenua, 839
mushroom sciarid fly, 839
soil moisture,
emergence, 465
insect pest, 465
life cycle, 465
Sminthurus viridis, 465
Solenopsis,
Popillia japonica, 2009
predation, 2009
secondary pest outbreak, 2009
Solenopsis invicta,
interspecific competition, 614
Monomorium chinense, 614
Pheidole fervens, 614
Solenopsis invicta virus-1, 614
Solenopsis invicta virus-1,
interspecific competition, 614
Monomorium chinense, 614
Pheidole fervens, 614
Solenopsis invicta, 614
solitary bees,
bee dispersal, 750
bee retention, 750
Osmia lignaria, 750
soybean aphid, 115, 1357, 2068
aphid-bioassay, 1357
Aphis glycines, 1357
EPG, 2068
feeding behavior, 2068
herbivore-induced plant volatile, 115
insecticide susceptibility, 1357
lacewing, 115
methyl salicylate, 115
resistance, 2068
syrphid fly, 115
Spalangia,
Musca domestica, 1108
Muscidifurax, 1108
population dynamics, 1108
Stomoxys calcitrans, 1108
Spathius agrili,
Agrilus planipennis, 379
biocontrol, 379
parasitoid rearing, 379
spatial distribution,
apterae, 1116
extraction, 1116
poplar petiolegall aphid, 1116
sampling, 1116
species displacement,
biological invasion, 1771
insecticide susceptibility, 1771
Liriomyza sativae, 1771
Liriomyza trifolii, 1771
species identification, 1280, 1720
DNA barcoding, 1720
internal transcribed spacer, 1720
lettuce, 1280
mitochondrial DNA, 1280
multiplex polymerase chain reaction, 1280
Plusiinae, 1280
Psocoptera, 1720
spinosad,
Anastrepha ludens, 452
fruit fly, 452
malathion, 452
traps, 452
spinosad bait,
neonicotinoid insecticides, 194
reproductively mature flies, 194

Rhagoletis indifferens, 194
sucrose, 194
spiroplasmas,
corn leafhopper, 1793
corn stunt, 1793
Dalbulus maidis, 1793
SPLAT,
citrus leafminer, 540
sex pheromone, 540
(Z,Z)-7,11-hexadecadienal, 540
(Z,Z,E)-7,11,13-hexadecatrienal, 540
Spodoptera exigua, 1229, 1343
carbohydrates, 1229
generating susceptible strain, 1343
insecticide resistance, 1343
lipids, 1229
methoxyfenozide, 1229
proteins, 1229
resistance management, 1343
starch,
developmental period, 2087
flour, 2087
population growth, 2087
Tribolium castaneum, 2087
starvation,
flight initiation, 2078
mating status, 2078
presence or absence of food, 2078
Tribolium castaneum, 2078
state-space model,
invasive insect species, 764
Kalman filter, 764
pests, 764
statistical power,
causal inference, 331
ecoinformatics, 331
economic injury level, 331
observational studies, 331
Steinernema,
Beauveria bassiana, 14
carbaryl, 14
Curculio caryae, 14
cypermethrin, 14
Steinernema carpocapsae,
Heterorhabditis, 47
peach, 47
Prunus persicae, 47
Synanthedon pictipes, 47
stereochemistry,
chrysanthemyl 2-methylbutanoate, 823
field bioassay, 823
lavandulyl 2-methylbutanoate, 823
Madeira mealybug, 823
sterile insect technique, 266, 420, 814, 1301, 1999
codling moth, 814
elytral color polymorphism, 420
Epiphyas postvittana, 1301, 1999
eradication, 1301, 1999
inherited sterility, 1999
irradiation, 1999
larval development, 266
long-distance transportation, 814
mass-marking method, 420
mating compatibility, 420
mosquito control, 266
pupation rate, 266
radiation, 1301
survival rate, 266
Tortricidae, 1301
sterile male releases,
Bactrocera dorsalis, 1969
invasive species, 1969
mating competitiveness, 1969
sticky gel,
hot melt pressure sensitive adhesive, 909
trap design, 909
western cherry fruit fly, 909
yellow color, 909
stink bug,
cold storage, 1833
conservational biological control, 1833
Gryon japonicum, 1833
Ooencyrtus nezarae, 1833
Stomoxys calcitrans, 250, 1108
Musca domestica, 1108
Muscidifurax, 1108
population dynamics, 1108
pyrethroid, 250
Spalangia, 1108
voltage-gated sodium channel, 250
storage,
aeration, 702
insects, 702
modeling, 702
rice, 702
stored grain,
integrated pest management, 325
nonchemical control, 325
stored grain beetles,
insect control, 308
ozone, 308
psocids, 308
stored wheat, 308
stored product, 1430, 1436, 1455
development, 1436
insect vision, 1430
LED, 1430
life table, 1436
oxygen, 1455
phosphine fumigation, 1455
pitfall trapping, 1430
postharvest pest control, 1455
psocid, 1436
temperature, 1436
stored wheat,
insect control, 308
ozone, 308
psocids, 308
stored grain beetles, 308
sublethal effect, 833, 970, 1942
Anoplophora glabripennis, 1942
Bemisia tabaci, 833
Calepitrimerus vitis, 970
fecundity, 1942
imidacloprid, 833
integrated pest management, 970
invasive species, 1942
neonicotinoid, 1942
predatory mites, 970
toxicity, 833, 970
substrate type,
Coptotermes formosanus, 2025
delayed action, 2025
nonrepellent, 2025
termiticide, 2025
subterranean termite, 1024, 1339, 1687
feeding rates, 1024
Heterotermes aureus, 1339
integrated pest management, 1687
invasive species, 1687
monitoring station, 1024
monitors, 1339
seasonal trends, 1024
succession ecology, 1687
trapping, 1339
succession ecology,
integrated pest management, 1687
invasive species, 1687
subterranean termite, 1687
sucrose,
neonicotinoid insecticides, 194
reproductively mature flies, 194
Rhagoletis indifferens, 194
spinosad bait, 194
sugarcane,
burnt cane, 155
green cane, 155
lesser cornstalk borer, 155
tillage, 155
Suleima helianthana,
Neotephritis finalis, 1236
planting date, 1236
sunflower, 1236
sunflower bud moth, 1236
sum of effective temperatures,
base temperature, 1870
day degrees, 1870
lower developmental threshold, 1870
rate of development, 1870
sunflower,
Neotephritis finalis, 1236
planting date, 1236
Suleima helianthana, 1236
sunflower bud moth, 1236
sunflower bud moth,
Neotephritis finalis, 1236
planting date, 1236
Suleima helianthana, 1236
sunflower, 1236
supercooling point,
Acari, 1423
Acaridae, 1423
disinfestation, 1423
nonchemical control, 1423

surface treatment,
insect growth regulator, 1765
Liposcelis, 1765
Psocoptera, 1765
pyriproxyfen, 1765
surveillance,
dairy, 1093
house fly, 1093
IPM, 1093
monitoring, 1093
survey,
disturbance, 879
edge, 879
exotic species, 879
trapping, 879
survival, 509, 532
degree-days, 532
development, 532
fecundity, 509
longevity, 509
navel orangeworm, 532
phenology, 509
pistachios, 532
rearing, 509
survival rate,
larval development, 266
mosquito control, 266
pupation rate, 266
sterile insect technique, 266
survivorship,
Aethina tumida, 753
diet, 753
oviposition, 753
temperature, 753
sweep net,
cotton, 685
plant bug, 685
sampling, 685
sweetpotato,
host plant resistance, 143
sweetpotato flea beetle, 143
wireworm, 143
sweetpotato flea beetle,
host plant resistance, 143
sweetpotato, 143
wireworm, 143
sweetpotato whitefly,
Bemisia tabaci, 548
Euphorbia pulcherrima, 548
poinsettia, 548
Synanthedon pictipes,
Heterorhabditis, 47
peach, 47
Prunus persicae, 47
Steinernema carpocapsae, 47
synchronization,
Dasineura mali, 1640
parasitism, 1640
phenology, 1640
Platygaster demades, 1640
synergism,
alcohol, 1680
cockroach, 1680
oil, 1680
toxicity, 1680
synergist,
chemical control, 1349
esterases, 1349
Mediterranean fruit fly, 1349
resistance management, 1349
synthetic oil,
Alpinia conchigera, 1445
contact toxicity, 1445
Curcuma zedoaria, 1445
Zingiber zerumbet, 1445
syrphid fly,
herbivore-induced plant volatile, 115
lacewing, 115
methyl salicylate, 115
soybean aphid, 115
systemic insecticides, 773, 1272
beneficial insects, 773
egg parasitoids, 773
fipronil, 1272
imidacloprid, 1272
laboratory bioassay, 773
Nantucket pine tip moth, 1272
pales weevil, 1272
predators, 773

T

T. canadensis,
fertility, 288
hemlock woolly adelgid, 288
T. heterophylla, 288
T. mertensiana, 288
T. heterophylla,
fertility, 288
hemlock woolly adelgid, 288
T. canadensis, 288
T. mertensiana, 288
T. mertensiana,
fertility, 288
hemlock woolly adelgid, 288
T. canadensis, 288
T. heterophylla, 288
target mutation,
Cydia molesta, 636
insecticide resistance, 636
mechanism, 636
monitoring, 636
temperature, 753, 1436
Aethina tumida, 753
development, 1436
diet, 753
life table, 1436
oviposition, 753
psocid, 1436
stored products, 1436
survivorship, 753
temperature threshold,
biological control, 799
fecundity, 799
intrinsic rate of increase, 799
population growth, 799
Tenebrio molitor,
bioassay, 2095
Ixodes scapularis, 2095
mealworms, 2095
Metarhizium brunneum, 2095
Tephritidae,
albedo, 54
flavedo, 54
host status, 54
Rutaceae, 54
termite, 425, 1502
antitermitic activity, 1502
Bacillus subtilis subsp. *virginiana*, 1502
biocontrol, 1502
cations, 425
foraging, 425
Reticulitermes flavipes, 425
soil, 425
termite feeding,
Klason lignin, 862
Odontotermes formosanus, 862
wood extractive, 862
wood hardness, 862
termite lipid content,
Coptotermes formosanus, 232
paper consumption, 232
Reticulitermes speratus, 232
termite survival, 232
termite survival,
Coptotermes formosanus, 232
paper consumption, 232
Reticulitermes speratus, 232
termite lipid content, 232
termiticide, 990, 2025
chlorantraniliprole, 990
Coptotermes formosanus, 2025
delayed action, 2025
indoxacarb, 990
nonrepellent, 2025
persistence, 990
R. flavipes, 990
substrate type, 2025
Tetranychidae,
biological control, 1496
integrated pest management, 1496
secondary pest outbreaks, 1496
tree nuts, 1496
Tetranychus cinnabarinus,
acaricidal activity, 375
Inula japonica, 375
Tetranychus urticae,
Deverra scoparia, 1220
essential oil, 1220
fecundity, 1220
toxicity, 1220
thermal requirements,
development rate, 1864
fecundity, 1864
population growth rate, 1864
Thrips tabaci,
Allium cepa, 1
Iris yellow spot virus, 1
management, 1
tillage,
burnt cane, 155
green cane, 155
lesser cornstalk borer, 155
sugarcane, 155

Tortricidae, 1301, 1514, 1706
barcodes, 1706
Epiphyas postvittana, 1301
eradication, 1301
host range, 1514
host specificity, 1514
internal transcribed spacer, 1706
invasive species, 1514
pest risk analysis, 1514
radiation, 1301
real-time polymerase chain reaction, 1706
sterile insect technique, 1301
toxicity, 833, 970, 1220, 1534, 1680
alcohol, 1680
Alpinia galanga, 1534
Bactrocera dorsalis, 1534
Bemisia tabaci, 833
Calepitrimerus vitis, 970
carboxylesterase, 1534
cockroach, 1680
Deverra scoparia, 1220
essential oil, 1220
fecundity, 1220
glutathione transferase, 1534
imidacloprid, 833
integrated pest management, 970
oil, 1680
predatory mites, 970
sublethal effect, 833
sublethal effects, 970
synergism, 1680
Tetranychus urticae, 1220
toxicity assay,
Bacillus thuringiensis, 1045
Diabrotica virgifera virgifera, 1045
MIR604, 1045
toxicity ranking,
Coptotermes formosanus, 1002
lethal contraction, 1002
lethal dose, 1002
Reticulitermes flavipes, 1002
toxin interaction,
clothianidin, 484
compensatory overconsumption, 484
hormesis, 484
hormoligosis, 484
transgenic Bt rice,
Folsomia candida, 1892
nontargets, 1892
risk assessment, 1892
transgenic corn,
blended seed refuge, 806
European corn borer, 806
feeding behavior, 806
resistance management, 806
transpiration,
Empoasca fabae, 1316
host plant expansion, 1316
photosynthesis, 1316
trap, 274, 452, 1018
Anastrepha ludens, 452
bed bug, 274
Cimex lectularius, 274
Drosophila, 1018
dry ice, 274
fruit fly, 452
malathion, 452
monitoring, 274
natural lure, 1018
spinosad, 452
trap color,
emerald ash borer, 901
funnel trap, 901
prism trap, 901
visual cues, 901
trap design, 909, 1258
attractant, 1258
Cerambycidae, 1258
hot melt pressure sensitive adhesive, 909
monitoring, 1258
sticky gel, 909
western cherry fruit fly, 909
woodborer, 1258
yellow color, 909
trapping, 879, 1339
disturbance, 879
edge, 879
exotic species, 879
Heterotermes aureus, 1339
monitors, 1339
subterranean termites, 1339
survey, 879
tree height,
Adelges tsugae, 1918
distribution, 1918
eastern hemlock, 1918
Tsuga canadensis, 1918
tree nuts,
biological control, 1496
integrated pest management, 1496
secondary pest outbreaks, 1496
Tetranychidae, 1496
Tribolium castaneum, 443, 1415, 2078, 2087
developmental period, 2087
flight initiation, 443, 2078
flour, 2087
food deprivation, 443
heat treatment, 1415
Liposcelis corrodens, 1415
Liposcelis decolor, 1415
mating status, 2078
population growth, 2087
presence or absence of food, 2078
rearing density, 443
red flour beetle, 443
Rhyzopertha dominica, 1415
starch, 2087
starvation, 2078
trichomes,
canola, 125
flea beetle feeding, 125
host plant resistance, 125
Phyllotreta spp., 125
Tsuga canadensis,
Adelges tsugae, 1918
distribution, 1918
eastern hemlock, 1918
tree height, 1918
turfgrass,
Blissus occiduus, 2073
Buchloe dactyloides, 2073
plant resistance, 2073
two-sex life table,
age-stage, 1525
Chilades pandava peripatria, 1525
Cycas revoluta, 1525
Cycas taitungensis, 1525

U

2-(undecyloxy)-ethanol,
aggregation pheromone, 1592
ethanol, 1592
Monochamus alternatus, 1592
α-pinene, 1592
urban entomology,
Arachnida, 601
Holocnemus pluchei, 601
pest control, 601

V

Vaccinium spp., 566, 571
Bactrocera cucurbitae, 566, 571
Bactrocera dorsalis, 566, 571
Bactrocera latifrons, 571
Ceratitis capitata, 566, 571
resistance, 566
Varroa destructor, 745, 1146, 1774
Apis mellifera, 745
asynchronous development, 1146
breeding, 745
carbon dioxide, 1774
environment, 1774
mating failure, 1146
mite resistance, 745
modified atmosphere, 1774
mortality, 1774
nonreproduction, 1146
resistance, 1146
vector management,
almond leaf scorch, 367
almonds, 367
Draeculacephala minerva, 367
sharpshooter, 367
vegetational diversity,
agroforestry systems, 164
Coffea arabica, 164
leaf-cutting ants, 164
vibrokinesis,
Agrilus planipennis, 81
auditory cues, 81
cryptic, 81
vibrotaxis, 81
vibrotaxis,
Agrilus planipennis, 81
auditory cues, 81

cryptic, 81
vibrokinesis, 81
vineyard,
bunch damage, 580
Cyprus, 580
insecticides, 580
Lobesia botrana, 580
virus,
bumble bee, 732
honey bee, 732
host expansion, 732
virus vectors,
aphids, 1167
barley yellow dwarf, 1167
cereal yellow dwarf, 1167
Rhopalosiphum padi, 1167
visual cues, 436, 901
chemical cues, 436
Coleoptera, 436
Cosmopolites sordidus, 436
Curculionidae, 436
emerald ash borer, 901
funnel trap, 901
prism trap, 901
trap color, 901
Vitacea polistiformis,
competitive attraction, 939
mating disruption, 939
pheromone dispenser, 939
volatiles,
1-octen-3-ol, 1633
borneol, 1633
menthol, 1633
repellency, 1633
voltage-gated sodium channel,
pyrethroid, 250
Stomoxys calcitrans, 250

W

water-sensitive spray cards,
insecticide spraying, 1138
integrated pest management, 1138
potato psyllids, 1138
row crops, 1138
western bean cutworm,
benefit-cost ratio, 1900
binomial sampling, 1900
resampling software, 1900
western cherry fruit fly,
hot melt pressure sensitive adhesive, 909
sticky gel, 909
trap design, 909
yellow color, 909
wheat, 1080, 1384, 1406
aphid biotype, 1080
Hessian fly, 1384
interaction, 1406
Mayetiola destructor, 1384
phytohormones, 1384
resistance, 1406
rice, 1384
Sitobion avenae, 1080
wheat curl mite, 1406
wheat resistance to aphid, 1080
wheat streak mosaic virus, 1406
wheat curl mite,
interaction, 1406
resistance, 1406
wheat, 1406
wheat streak mosaic virus, 1406
wheat resistance to aphid,
aphid biotype, 1080
Sitobion avenae, 1080
wheat, 1080
wheat streak mosaic virus,
interaction, 1406
resistance, 1406
wheat, 1406
wheat curl mite, 1406
white trap,
Aethina tumida, 26
in-hive traps, 26
out-hive traps, 26
polyvinyl chloride trap, 26
wireworm,
host plant resistance, 143
sweetpotato, 143
sweetpotato flea beetle, 143
within-plant distribution,
ants, 947
aphid, 947
Banana bunchy top virus, 947
sampling, 947
wood extractive,
Klason lignin, 862
Odontotermes formosanus, 862
termite feeding, 862
wood hardness, 862
wood hardness,
Klason lignin, 862
Odontotermes formosanus, 862
termite feeding, 862
wood extractive, 862
woodborer,
attractant, 1258
Cerambycidae, 1258
monitoring, 1258
trap design, 1258
woolly whiteflies,
Aleurothrixus floccosus, 1329
citrus, 1329
cyhalothrin, 1329
selecron, 1329

X

X-ray irradiation,
Bactrocera cucurbitae, 1509
Bactrocera latifrons, 1509
phytosanitary treatment, 1509
quarantine pest, 1509
Xylosandrus crassiusculus, 1960, 2017
bottle-traps, 2017
economic analysis, 1960
ethanol concentration, 2017
insecticide coverage, 1960
Oligonycus aceris, 1960
secondary pest outbreak, 1960
Xylosandrus germanus, 2017
Xylosandrus germanus, 1665, 2017
allyl isothiocyanate, 1665
ambrosia beetles, 1665
botanical insecticides, 1665
bottle-traps, 2017
essential oils, 1665
ethanol concentration, 2017
Xylosandrus crassiusculus, 2017

Y

yellow color, 909
hot melt pressure sensitive adhesive, 909
sticky gel, 909
trap design, 909
western cherry fruit fly, 909
yield loss,
economic threshold, 1550
field pea, 1550
nitrogen fertilizer, 1550
pea leaf weevil, 1550
Y-tube bioassay,
Adelphocoris suturalis, 1189
circadian rhythm, 1189

Z

zebra chip, 586, 1486
inoculation efficiency, 1486
insecticides, 586
IPM, 586
liberibacter, 1486
potato, 586, 1486
potato psyllid, 586, 1486
zebra chip disease,
liberibacter, 1783
potato, 1783
potato psyllid, 1783
psyllid density, 1783
Z-3-hexenol,
ash canopy, 173
green prism traps, 173
monitoring, 173
phoebe oil, 173
Zingiber zerumbet,
Alpinia conchigera, 1445
contact toxicity, 1445
Curcuma zedoaria, 1445
synthetic oil, 1445
(Z,Z)-7,11-hexadecadienal,
citrus leafminer, 540
sex pheromone, 540
SPLAT, 540
(Z,Z)-7,11-hexadecadienal, 540
(Z,Z,*E*)-7,11,13-hexadecatrienal, 540
(Z,Z,*E*)-7,11,13-hexadecatrienal,
citrus leafminer, 540
sex pheromone, 540
SPLAT, 540
(Z,Z)-7,11-hexadecadienal, 540
(Z,Z,*E*)-7,11,13-hexadecatrienal, 540